

On localized structures in 2d and 3d models comprising Yang-Mills fields

I.L. Bogolubsky (JINR),
A.A. Bogolubskaya (JINR)

Symposium on Theoretical and
Mathematical Physics,
St. Petersburg,
July 8-13, 2011

Abstract: We discuss the possibility of soliton existence in 2D and 3D SU(2) gluodynamics in Lorentz gauge. Hamiltonians in terms of radial functions are presented. We are looking for localized in space YM field distributions which provide local minima to these hamiltonians. Such nontopological solitons if exist may be relevant to extended gluonic strings in mesons (in 2D) and glueball states (in 3D). Finally separation of variables and Hamiltonian density are presented for 3D SU2-Higgs EW model.

Quark-antiquark with gluonic string

The famous action density distribution between two static colour sources [G.S. Bali, K. Schilling, C. Schlichter '95].





Figure 1: Structure of mesons

Introduction

- Until now there is no satisfactory theoretical description of extended string connecting quark and antiquark in mesons.
- Study of 2D solitons can clarify this issue.
- For now nobody proposed adequate ansatz for description of $2D$ Yang-Mills solitons.
- For 3D case only the simplest one-term ansatz has been studied, for it $\partial_\mu A_\mu = 0$ is valid.
- Generic 3-term ansatz requires detailed study, for it $\partial_\mu A_\mu = 0$ is not automatically satisfied.
- 3D YM solitons if exist could be viewed as classical glueballs.
- In previous studies of Yang-Mills solitons specifics of Yang-Mills fields as gauge ones has been never taken into account.
- Non-perturbative effects in Salam-Weinberg EW theory are not sufficiently taken into account for now.

Ansatz for Yang-Mills in D=2

- Consider the vector $SU(2)$ Yang-Mills field $A_\mu^a(x^\nu)$,

$$\mathcal{L} = -\frac{1}{4}(F_{\mu\nu}^a)^2,$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\varepsilon^{abc} A_\mu^b A_\nu^c,$$

$$D = 2, \quad \mu, \nu = 0, 1, 2, \quad a, b, c = 1, 2, 3, \quad g - \text{const.}$$

- We look for stationary solutions and use the following ansatz:

$$A_0^a = 0,$$

$$gA_i^a = \delta_{a3} \varepsilon_{iak} x_k \frac{1}{R^2} s(R) +$$

$$+ (\delta_{a1} + \delta_{a2}) \left[(\delta_{ia} R^2 - x_i x_a) \frac{b(R)}{R^3} + \frac{p(R) x_i x_a}{R^4} \right],$$

$$i, k = 1, 2 \quad R^2 = x^2 + y^2.$$

Hamiltonian density for $D=2$

No gauge fixing here.

> **H_YM_2D;**

$$\begin{aligned}
 & \frac{1}{2} \frac{s(R)^2 p(R)^2}{R^2 g^2} - \frac{b(R) p(R)}{R^3 g^2} - \frac{p(R) \left(\frac{d}{dR} b(R) \right)}{R^2 g^2} + \frac{1}{2} \frac{\left(\frac{d}{dR} b(R) \right)^2}{g^2} \\
 & + \frac{p(R)^2 s(R)}{R^3 g^2} + \frac{b(R) \left(\frac{d}{dR} b(R) \right)}{R g^2} + \frac{1}{2} \frac{b(R)^2}{R^2 g^2} + \frac{1}{2} \frac{\left(\frac{d}{dR} s(R) \right)^2}{g^2} \\
 & + \frac{1}{2} \frac{b(R)^2 p(R)^2}{R^2 g^2} + \frac{1}{2} \frac{s(R)^2}{R^2 g^2} + \frac{s(R) \left(\frac{d}{dR} s(R) \right)}{R g^2} \\
 & + \frac{\left(\frac{d}{dR} s(R) \right) b(R) p(R)}{R g^2} - \frac{\left(\frac{d}{dR} b(R) \right) s(R) p(R)}{R g^2} + \frac{1}{2} \frac{p(R)^2}{R^4 g^2}
 \end{aligned}$$

Maple output 1: Hamiltonian density, $D=2$.

Yang-Mills in $D = 3$ (1)

- Consider the vector $SU(2)$ Yang-Mills field $A_\mu^a(x^\nu)$,

$$\mathcal{L} = -\frac{1}{4}(F_{\mu\nu}^a)^2,$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\varepsilon^{abc} A_\mu^b A_\nu^c,$$

$$D = 3, \quad \mu, \nu = 0, 1, 2, 3 \quad a, b, c = 1, 2, 3, \quad g - \text{const.}$$

- Generic ansatz for $D = 3$ YM solitons:

$$A_0^a = \frac{x^a}{R} q(R),$$

$$gA_i^a = \varepsilon_{iak} \frac{x_k}{R^2} s(R) + \frac{b(R)}{R^3} (\delta_{ia} R^2 - x_i x_a) + \frac{p(R) x_i x_a}{R^4};$$

$$i, k = 1, 2, 3, \quad R^2 = x^2 + y^2 + z^2.$$

Yang-Mills in $D = 3$ (2)

> **H_YM_3D;**

$$\begin{aligned}
 & \frac{\left(\frac{d}{dR} b(R)\right)^2}{R^2} + \frac{\left(\frac{d}{dR} s(R)\right)^2}{R^2} + \frac{2 s(R)^3}{R^4} + \frac{2 s(R) b(R)^2}{R^4} \\
 & - \frac{2 \left(\frac{d}{dR} b(R)\right) s(R) p(R)}{R^4} + \frac{1}{2} \frac{b(R)^4}{R^4} + \frac{2 b(R) p(R) \left(\frac{d}{dR} s(R)\right)}{R^4} \\
 & + \frac{2 s(R)^2}{R^4} - \frac{2 p(R) \left(\frac{d}{dR} b(R)\right)}{R^4} + \frac{b(R)^2 s(R)^2}{R^4} + \frac{1}{2} \frac{s(R)^4}{R^4} + \frac{p(R)^2}{R^6} \\
 & + \frac{s(R)^2 p(R)^2}{R^6} + \frac{b(R)^2 p(R)^2}{R^6} + \frac{2 p(R)^2 s(R)}{R^6} + \frac{3}{4} \frac{q(R)^2 \left(\frac{d}{dR} p(R)\right)^2}{R^2} \\
 & + \frac{1}{4} q(R)^2 + \frac{3}{2} \left(\frac{d}{dR} q(R)\right)^2 + \frac{3 (s(R) + 1)^2 q(R)^2}{R^2}
 \end{aligned}$$

>

Maple output 2: Hamiltonian density, $D=3$, no gauge fixing.

Apply Lorentz gauge $\partial_\mu A_\mu = 0$

Now apply Lorentz gauge $\partial_\mu A_\mu = 0$.

For $D=2$ Hamiltonian density takes the form:

$$\mathcal{H}_{sol} = \frac{1}{2g^2} \left[\left(\frac{ds}{dR} + \frac{s}{R} + \frac{p}{R^3} \frac{dp}{dR} \right)^2 + \frac{1}{R^2} \left(\frac{d^2p}{dR^2} - \frac{p}{R} \left(s + \frac{1}{R} \right) \right)^2 \right] \quad (1)$$

For $D=3$ Hamiltonian density reads:

$$\begin{aligned} \mathcal{H}_{sol} = & \frac{1}{g^2} \left\{ \frac{1}{32 R^4} \left[\left(\frac{dp}{dR} \right)^2 + 8s + 4s^2 \right]^2 + \right. \\ & \left[\frac{p(s+1)}{R^3} - \frac{1}{2R} \frac{d^2p}{dR^2} \right]^2 + \left[\frac{1}{R} \frac{ds}{dR} + \frac{1}{2R^3} \frac{dp}{dR} p \right]^2 + \\ & \left. \left[\frac{3}{4} \left(\frac{dp}{dR} \right)^2 + \frac{1}{4} + \frac{3(s+1)^2}{R^2} \right] q^2 + \frac{3}{2} \left(\frac{dq}{dR} \right)^2 \right\} \quad (2) \end{aligned}$$

\Rightarrow Numerical search for localized solutions is in progress. We plan to start with Monte-Carlo simulations.

No-Go Theorems, Coleman & Co. (1)

Coleman's study: let $A_\mu^a(x)$ - classical localized solution. Make transformations

$$\begin{aligned} A_0^a(x_k; \sigma, \lambda) &= \sigma \lambda A_0^a(\lambda x_k), \\ A_i^a(x_k; \sigma, \lambda) &= \lambda A_i^a(\lambda x_k). \end{aligned} \tag{1}$$

Denote

$$\begin{aligned} H_1 &= \frac{1}{2} \int d^{\mathcal{D}}x (F_{0i}^a)^2 \\ &= \frac{1}{2} \int d^{\mathcal{D}}x (\partial_i A_0^a + e c^{abc} A_0^b A_i^c)^2, \end{aligned} \tag{2}$$

$$\begin{aligned} H_2 &= \frac{1}{2} \int d^{\mathcal{D}}x (F_{ij}^a)^2 \\ &= \frac{1}{4} \int d^{\mathcal{D}}x (\partial_j A_i^a + e c^{abc} A_i^b A_j^c)^2. \end{aligned} \tag{3}$$

Then under transformation (1)

$$H(\sigma, \lambda) = \sigma^2 \lambda^{(4-\mathcal{D})} H_1 + \lambda^{(4-\mathcal{D})} H_2 .$$

No-Go Theorems, Coleman & Co. (2)

Requiring stationarity:

$$\frac{\partial H}{\partial \sigma} = 0, \quad \frac{\partial H}{\partial \lambda} = 0 \quad \text{at} \quad \sigma = 1, \quad \lambda = 1,$$

Coleman has found for $D \neq 4$: $H_1 = H_2 = 0$.

For $D \neq 4$ from here: $F_{\mu\nu}^a = 0$, *Q.E.D.*

Coleman's conclusion was:

“There are no classical glueballs”.

⇒ Thus, Coleman has shown that there are no minima of Hamiltonian in extended space of variables, corresponding to non-fixed gauge fields and including nonphysical degrees of freedom. E.g. fixing the Lorentz gauge, we get the physical space of dynamical variables, whose dimensionality is less than that of extended space of gauge field **without gauge fixing**.

⇒ In such physical space the existence of minima is not forbidden. Hence we can hope that $3D$ *YM* solitons exist.

EW SU2-Higgs model

(1)

- Lagrangian density:

$$\mathcal{L} = |\mathcal{D}_\mu \varphi|^2 - \frac{1}{4}(F_{\mu\nu}^a)^2,$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g\varepsilon^{abc} A_\mu^b A_\nu^c,$$

$$\varphi = \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix}, \quad \mathcal{D}_\mu \varphi = \left(\partial_\mu - \frac{ig}{2} \tau_a A_\mu^a \right) \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix}.$$

Here φ is isospinor of SU(2) group,
 $D = 3$, $\mu, \nu = 0, 1, 2, 3$; $a, b, c = 1, 2, 3$; $g - const.$

- Isospinor φ is represented by four real values ϕ_α , ($\alpha = 0, 1, 2, 3$):

$$\varphi_1 = \frac{\phi_1 + i\phi_2}{\sqrt{2}},$$

$$\varphi_2 = \frac{\phi_0 + i\phi_3}{\sqrt{2}},$$

such that $\phi_0^2 + \phi_1^2 + \phi_2^2 + \phi_3^2 = 1$.

i.e. unit 4-vector ϕ_α takes values on unit sphere S^3 .

EW SU2-Higgs model (2)

- Localized 3D configurations of quasi-Higgs field φ ,
 $\varphi(\infty) = \varphi_0$,

define maps $R_{comp}^3 \rightarrow S^3$ or equivalently
 $S^3 \rightarrow S^3$.

Hence φ -configurations with nontrivial topological indices (mapping degrees Q_{top}) are possible.

- Consider the case $Q_{top} = 1$ and try the following **ansatz** for EW SU2-Higgs model:

$$A_0^a = \frac{x^a}{R} q(R),$$

$$gA_i^a = \varepsilon_{iak} \frac{x_k}{R^2} s(R) + \frac{1}{2} \frac{dp(R)}{dR} \frac{1}{R^3} (\delta_{ia} R^2 - x_i x_a) +$$

$$+ \frac{p(R) x_i x_a}{R^4},$$

$$\varphi = \begin{pmatrix} \varphi_1 \\ \varphi_2 \end{pmatrix} = \left[\sin \theta(R) \frac{\tau_a x_a}{R} + i \cos \theta(R) \right] \begin{bmatrix} 1 \\ 0 \end{bmatrix},$$

$$\tau_1 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \tau_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$\cos \theta(0) = -1, \quad \cos \theta(\infty) = 1.$$

EW SU2-Higgs model (3)

Also apply Lorentz gauge $\partial_\mu A_\mu = 0$.

For $D=3$ Hamiltonian density reads:

$$\begin{aligned} \mathcal{H}_{sol} = & \frac{1}{g^2} \left\{ \frac{1}{32 R^4} \left[\left(\frac{dp}{dR} \right)^2 + 8s + 4s^2 \right]^2 + \right. \\ & \left[\frac{p(s+1)}{R^3} - \frac{1}{2R} \frac{d^2 p}{dR^2} \right]^2 + \left[\frac{1}{R} \frac{ds}{dR} + \frac{1}{2R^3} \frac{dp}{dR} p \right]^2 + \\ & \left[\frac{3}{4} \frac{1}{R^2} \left(\frac{dp}{dR} \right)^2 + \frac{1}{4} + \frac{3(s+1)^2}{R^2} \right] q^2 + \frac{3}{2} \left(\frac{dq}{dR} \right)^2 + \\ & \left. \left(\frac{d\theta}{dR} \right)^2 + \frac{1}{R^2} \left[2 \sin^2(\theta)(s+1) + \frac{1}{2} \frac{dp}{dR} \sin(2\theta) + p \frac{d\theta}{dR} \right] \right\}. \end{aligned}$$

\Rightarrow Numerical search for localized solutions is in progress. We plan to start with Monte-Carlo simulations.

References

1. G.S. Bali, K. Schilling and C. Schlichter, Observing long color flux tubes in SU(2) lattice gauge theory, PRD 51 (1995) 5165; hep-lat/9409005.
2. I.L. Bogolubsky and A. A. Bogolubskaya, On Two-Field Solitons in 2 and 3 Dimensions, Proc. of Int. Conf. "Mathematical Modeling and Computational Physics(MMCP 2009)", No 2, Issue 2. (2010) 158.
3. S. Coleman, Classical lumps and their quantum descendants, Proc. of the 1975 Intern. School of Subnuclear Physics. Part A. Ed. by Zichichi (N. Y.—London: Plenum Press, 1976) 297.
4. S. Coleman, There are no classical glueballs, Commun. Math. Phys. 55 (1977) 113.
5. S. Deser, Absence of static EYM excitations in three dimensions, PLB 64 (1976) 463.
6. V.A. Rubakov, Classical Gauge Fields (Inst. of Nucl. Research, Moscow, 1998) (in Russian), parts 1, 2, 3.
7. I.L. Bogolubsky and A. A. Bogolubskaya, On 2D and 3D solitons in SU(2) gluodynamics, in proceedings of ICHEP2010, PoS(ICHEP 2010)086.
URL: http://pos.sissa.it/archive/conferences/120/086/ICHEP_2010_086.pdf