

# Abelian and nilpotent regular maps and dessins

Roman Nedela

Matej Bel University/Math. Inst. Slovak Academy of Sciences

Banská Bystrica

Saint Petersburg, October 2014

October 26, 2014

Original results on the nilpotent maps were done in collaboration with **M. Conder, S.F. Du, A. Malnič and M. Škoviera.**

Original results on the nilpotent dessins come from the thesis of my former PhD. student **Naer Wang** defended on May, 29, 2014 and from subsequent papers in progress by **Kan Hu, Naer Wang** and N.

# Maps and Combinatorial Maps

**Topological map** A 2-cell decomposition of a surface = 2-cell embedding of a connected graph into a surface,

**Surfaces** preferably compact connected and orientable and their universal covers,

**Combinatorial Map** Equivalence classes of topological maps up to isotopy can be described by triples  $(D; R, L)$

$D$  a set of darts,  $L$  is an (dart-reversing) involution,  $R$  rotation system,

**Category of Ormaps** Combinatorial maps and their homomorphisms,

**Vertices, Edges and Faces** Orbits of  $R$ ,  $L$ , and  $RL$ ,

# Regular maps

**Monodromy group** is the group  $\langle R, L \rangle$ , by definition it is transitive on the set  $D$  of darts,

**Automorphism group** permutations in  $Sym(D)$  centralising the monodromy group form the automorphism group  $Aut(M)$  of a map  $M$ ,

**semiregularity**  $Aut(M)$  is semiregular on  $D$ ;

**regularity** if  $Aut(M)$  is regular on  $D$ , then  $M$  is called regular;  
in a regular map  $Aut(M) \cong Mon(M)$  and  
 $|D| = |Aut(M)| = |Mon(M)|$ ;

# Abelian and Nilpotent maps

A regular map with an (abelian) nilpotent automorphism group will be called an **(abelian) nilpotent map**,

# Abelian and Nilpotent maps

A regular map with an (abelian) nilpotent automorphism group will be called an **(abelian) nilpotent map**,

A (abelian) nilpotent map can be view as a triple  $(G; x, y)$ , where  $G = \langle x, y \rangle$  is an (abelian) nilpotent group,  $y^2 = 1$ ,

# Abelian and Nilpotent maps

A regular map with an (abelian) nilpotent automorphism group will be called an **(abelian) nilpotent map**,

A (abelian) nilpotent map can be view as a triple  $(G; x, y)$ , where  $G = \langle x, y \rangle$  is an (abelian) nilpotent group,  $y^2 = 1$ ,

Two such maps (triples) are **equivalent** if there the assignment  $x_1 \mapsto x_2$  and  $y_1 \mapsto y_2$  extends to a group isomorphism,

# Problem(s) considered

General problem: Characterisation (classification) of nilpotent maps.

# Problem(s) considered

General problem: Characterisation (classification) of nilpotent maps.

More realistic problems: Investigate invariants: nilpotency **class, type, structure of underlying graphs, chirality/reflexibility, classification of particular subclasses,**

# Problem(s) considered

General problem: Characterisation (classification) of nilpotent maps.

More realistic problems: Investigate invariants: nilpotency **class, type, structure of underlying graphs, chirality/reflexibility, classification of particular subclasses,**

In general, we want to understand what are the most important invariants of nilpotent maps,

## Problem(s) considered

General problem: Characterisation (classification) of nilpotent maps.

More realistic problems: Investigate invariants: nilpotency **class, type, structure of underlying graphs, chirality/reflexibility, classification of particular subclasses,**

In general, we want to understand what are the most important invariants of nilpotent maps,

Since **non-abelian nilpotent maps are bipartite,** some of the ideas naturally generalise to **nilpotent regular hypermaps and dessins.**

# Abelian regular maps, an easy exercise

## Theorem

*A folklore result: Abelian regular maps are the semistars, balanced regular embeddings of bouquets of loops and embeddings of  $n$ -dipoles with exponent 1.*

An argument: the group is either cyclic, or  $Z_2 \times Z_n$ , check how one can generate them by an involution and a non-involution.

# Examples of nilpotent maps

- ① Cycles of length  $n$ ,  $n = 2^m$ , and their duals, of nilpotency class  $m$

# Examples of nilpotent maps

- ① Cycles of length  $n$ ,  $n = 2^m$ , and their duals, of nilpotency class  $m$
- ② Regular embeddings of cycles of length  $n = 2^m$  and multiplicity  $2^k$

# Examples of nilpotent maps

- ① Cycles of length  $n$ ,  $n = 2^m$ , and their duals, of nilpotency class  $m$
- ② Regular embeddings of cycles of length  $n = 2^m$  and multiplicity  $2^k$
- ③ Du, Jones, Kwak, Nedela, Skoviera (2007, 2010, 2013): Regular embeddings of  $K_{n,n}$ , where  $n = 2^m$ , there are  $2^{m-2}$  of them with the metacyclic colour preserving group and four with the non-metacyclic subgroup.

# Examples of nilpotent maps

- ① Cycles of length  $n$ ,  $n = 2^m$ , and their duals, of nilpotency class  $m$
- ② Regular embeddings of cycles of length  $n = 2^m$  and multiplicity  $2^k$
- ③ Du, Jones, Kwak, Nedela, Skoviera (2007, 2010, 2013): Regular embeddings of  $K_{n,n}$ , where  $n = 2^m$ , there are  $2^{m-2}$  of them with the metacyclic colour preserving group and four with the non-metacyclic subgroup.
- ④ Catalano, Conder, Du, Kwon, N., Wilson (2011): Regular embeddings of  $Q_n$ ,  $n = 2^m$ , the number of them grows approximately as  $2^n$

# Examples of nilpotent maps

- ① Cycles of length  $n$ ,  $n = 2^m$ , and their duals, of nilpotency class  $m$
- ② Regular embeddings of cycles of length  $n = 2^m$  and multiplicity  $2^k$
- ③ Du, Jones, Kwak, Nedela, Skoviera (2007, 2010, 2013): Regular embeddings of  $K_{n,n}$ , where  $n = 2^m$ , there are  $2^{m-2}$  of them with the metacyclic colour preserving group and four with the non-metacyclic subgroup.
- ④ Catalano, Conder, Du, Kwon, N., Wilson (2011): Regular embeddings of  $Q_n$ ,  $n = 2^m$ , the number of them grows approximately as  $2^n$

Conclusion: **Nilpotent maps form a complex family of maps**

# Basic facts on the nilpotent maps

## Theorem (Malnic, N., Škoviera)

*A nilpotent map  $(G; x, y)$  decomposes into  $S$ . Wilson's product:  $G_2 \times S_{S_m}$ , where  $G_2 = \langle x_2, y_2 \rangle$  is the Sylow 2-group and  $S_{S_m}$  is an odd semistar,*

## Corollary

*The automorphism group of a nilpotent map with a simple underlying graph is a 2-group.*

# Nilpotent maps of class two

Theorem (Malnic, N., Škoviera EJC 2012)

*Every nilpotent map of class two belong to one of the two families defined by presentations:*

$$G_1(n) = \langle x, y \mid x^{2^n} = y^2 = 1, [x, y] = x^{2^{n-1}} \rangle, n \geq 2$$

*and*

$$G_2(n) = \langle x, y, z \mid x^{2^n} = y^2 = z^2 = [x, z] = [y, z] = 1, z = [x, y] \rangle.$$

# Nilpotent maps of class three I

joint work: Ban, Du, Liu, N., Škoviera (2014+):

$$H_1(k) = \langle a, b \mid a^{2^k} = b^2 = 1, [a, b] = c, [c, a] = [c, b] = e, [e, a] = [e, b] = 1 \rangle, k \geq 1;$$

$$H_2(k) = \langle a, b \mid a^{2^k} = e, b^2 = 1, [a, b] = c, [c, a] = [c, b] = e, [e, a] = [e, b] = 1 \rangle, k \geq 1;$$

$$H_3(k) = \langle a, b \mid a^{2^k} = b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle, k \geq 2;$$

$$H_4(k) = \langle a, b \mid a^{2^k} = d, b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = 1, [d, a] = [d, b] = 1 \rangle, k \geq 2;$$

# Nilpotent maps of class three II

$$\begin{aligned}H_5(k) &= \langle a, b \mid a^{2^k} = b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = e, \\ &\quad [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle, k \geq 2; \\ H_6(k) &= \langle a, b \mid a^{2^k} = d, b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = e, \\ &\quad [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle, k \geq 2; \\ H_7(k) &= \langle a, b \mid a^{2^k} = e, b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = e, \\ &\quad [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle, k \geq 3; \\ H_8(2) &= \langle a, b \mid a^4 = de, b^2 = 1, [a, b] = c, [c, a] = d, [c, b] = e, \\ &\quad [d, a] = [d, b] = [e, a] = [e, b] = 1 \rangle.\end{aligned}$$

# Projection and lifting of nilpotent maps

Given a nilpotent map one can form a chain of quotients by factoring out central subgroups, and one can do it in the way that for each class  $c' < c$  a map of class  $c'$  is included!

# Projection and lifting of nilpotent maps

Given a nilpotent map one can form a chain of quotients by factoring out central subgroups, and one can do it in the way that for each class  $c' < c$  a map of class  $c'$  is included!

More generally, nilpotent maps of bounded class are closed under taking quotients

# Projection and lifting of nilpotent maps

Given a nilpotent map one can form a chain of quotients by factoring out central subgroups, and one can do it in the way that for each class  $c' < c$  a map of class  $c'$  is included!

More generally, nilpotent maps of bounded class are closed under taking quotients

It can be used firstly to lift some properties of maps of smaller class, secondly to lift classification statements.

# Projection and lifting of nilpotent maps

Given a nilpotent map one can form a chain of quotients by factoring out central subgroups, and one can do it in the way that for each class  $c' < c$  a map of class  $c'$  is included!

More generally, nilpotent maps of bounded class are closed under taking quotients

It can be used firstly to lift some properties of maps of smaller class, secondly to lift classification statements.

Example: The underlying graphs of maps of class 2 are 4-cycles with multiple edges, or dipoles. Consequence: Non-abelian nilpotent maps are **bipartite!**

# Summary of basic properties

## Theorem

*The family of nilpotent maps of bounded class is:*

- (a) closed under Wilson's products,*
- (b) closed under taking quotients,*
- (c) admits lower central series intersecting all the families of smaller classes.*

## Universal map of class $c$

$\Gamma = C * C_2 \cong \langle x, y | y^2 = 1 \rangle$ , (the universal map)

Set  $\Gamma_1 = [\Gamma, \Gamma]$  and in general  $\Gamma_{n+1} = [\Gamma, \Gamma_n]$ .

Then the universal map of class  $c$  is

$$H(c) = \Gamma / \Gamma_c \cong \langle \bar{x}, \bar{y} \rangle,$$

## Universal map of class $c$

$\Gamma = C * C_2 \cong \langle x, y | y^2 = 1 \rangle$ , (the universal map)

Set  $\Gamma_1 = [\Gamma, \Gamma]$  and in general  $\Gamma_{n+1} = [\Gamma, \Gamma_n]$ .

Then the universal map of class  $c$  is

$$H(c) = \Gamma / \Gamma_c \cong \langle \bar{x}, \bar{y} \rangle,$$

it is an infinite group, but for small  $c \leq 3$  we observed that nilpotent maps tend to have parallel edges.

# Universal map of class $c$

$\Gamma = C * C_2 \cong \langle x, y | y^2 = 1 \rangle$ , (the universal map)

Set  $\Gamma_1 = [\Gamma, \Gamma]$  and in general  $\Gamma_{n+1} = [\Gamma, \Gamma_n]$ .

Then the universal map of class  $c$  is

$$H(c) = \Gamma / \Gamma_c \cong \langle \bar{x}, \bar{y} \rangle,$$

it is an infinite group, but for small  $c \leq 3$  we observed that nilpotent maps tend to have parallel edges.

**Conjecture:** there are just finitely many nilpotent maps of bounded class without parallel edges.

# The number of vertices of the universal map?

Group theoretical formulation:

Determine the size of  $U(c) = H(c)/\text{core}(\bar{x})$ .

# The number of vertices of the universal map?

Group theoretical formulation:

Determine the size of  $U(c) = H(c)/\text{core}(\bar{x})$ .

Surprise:  $U(c)$  is finite!!!

A recent result by Conder, Du, N., Škoviera: we determined a precise formula for the size of  $|U(c)|$ , its number of vertices, its type and genus.

# Three integer sequences

Set  $h_1 = 0$ , and then for  $n \geq 2$  take

$$h_n = \frac{1}{n} \sum_{d|n} \mu(n/d) \left( \sum_{0 \leq i \leq d/3} (-1)^i \frac{d}{d-2i} \binom{d-2i}{i} 2^{d-3i} \right),$$

$$g_n = h_1 + h_2 + \cdots + h_n = \sum_{j=1}^n h_j,$$

$$f_n = g_1 + g_2 + \cdots + g_n = \sum_{i=1}^n g_i.$$

As usual,  $\mu(m)$  denotes the Möbius function.

# The number of vertices of the universal map of class $c$

## Theorem (Labute77)

*The rank of the factor group  $\Gamma_{n-1}/\Gamma_n$  is  $g_n$ , for all  $n \geq 2$ .*

## Theorem (Conder, Du, N., Škoviera, 2014...)

*For every integer  $c \geq 1$  there exists a unique nilpotent regular map  $\mathcal{U}_c$  of class  $c$  with simple underlying graph, having  $2^{1+f_c}$  vertices, and type  $\{2^c, 2^{c-1}\}_{2^c}$ .*

*Furthermore, this map is 'universal', in the sense that every nilpotent regular map of class at most  $c$  with simple underlying graph is a quotient of  $\mathcal{U}_c$ .*

## Generation of simple universal maps of given class

Just add relation  $x^{c-1} = 1$  to the presentation of  $H(c)$ , since  $\text{core}(x) = \langle x^{c-1} \rangle$  and identify all the normal subgroups.

Example: There are 21 nilpotent maps of class  $c$ ,  $2 \leq c \leq 4$ , with a simple underlying graph.

## Generation of simple universal maps of given class

Just add relation  $x^{c-1} = 1$  to the presentation of  $H(c)$ , since  $\text{core}(x) = \langle x^{c-1} \rangle$  and identify all the normal subgroups.

Example: There are 21 nilpotent maps of class  $c$ ,  $2 \leq c \leq 4$ , with a simple underlying graph.

$$\text{M2.8.1} \quad [x^2, y^2, (xy)^4],$$

$$\text{M3.64.1} \quad [x^4, y^2, x^{-1}yx^{-2}yxyx^2y, (x^{-1}y)^4(xy)^4],$$

$$\text{M3.32.1} \quad [x^4, y^2, (x^{-1}y)^4, (xyx^{-1}y)^2]$$

$$\text{M3.32.2} \quad [x^4, y^2, x^{-1}yx^{-1}yxyxy]$$

$$\text{M3.16.1} \quad [x^2, y^2, (xy)^8]$$

## Generation of simple universal maps of given class

Just add relation  $x^{c-1} = 1$  to the presentation of  $H(c)$ , since  $\text{core}(x) = \langle x^{c-1} \rangle$  and identify all the normal subgroups.

Example: There are 21 nilpotent maps of class  $c$ ,  $2 \leq c \leq 4$ , with a simple underlying graph.

$$\text{M2.8.1} \quad [x^2, y^2, (xy)^4],$$

$$\text{M3.64.1} \quad [x^4, y^2, x^{-1}yx^{-2}yxyx^2y, (x^{-1}y)^4(xy)^4],$$

$$\text{M3.32.1} \quad [x^4, y^2, (x^{-1}y)^4, (xyx^{-1}y)^2]$$

$$\text{M3.32.2} \quad [x^4, y^2, x^{-1}yx^{-1}yxyxy]$$

$$\text{M3.16.1} \quad [x^2, y^2, (xy)^8]$$

**Conclusion: Nilpotent maps of given class can be sorted into families characterised by the smallest members which have no parallel edges!**

## Note on the sequences $h_n$ and $g_n$

First values of  $h_n$  are

0, 1, 1, 1, 2, 2, 4, 5, 8, 11, 18, 25, 40, 58, 90, 135, 210, 316, ...

## Note on the sequences $h_n$ and $g_n$

First values of  $h_n$  are

0, 1, 1, 1, 2, 2, 4, 5, 8, 11, 18, 25, 40, 58, 90, 135, 210, 316, ...

Sloane on-line Encyclopedia of integer sequences:

“A006206 Number of aperiodic binary necklaces of length  $n$  with no subsequence 00, excluding the necklace “0”. (Formerly M0317)

1, 1, 1, 1, 2, 2, 4, 5, 8, 11, 18, 25, 40, 58, 90, 135, 210, 316, 492, 750, 1164, 1791, 2786, 4305, 6710, 10420, 16264, 25350, 39650, 61967, 97108, “

## Note on the sequences $h_n$ and $g_n$

First values of  $h_n$  are

0, 1, 1, 1, 2, 2, 4, 5, 8, 11, 18, 25, 40, 58, 90, 135, 210, 316, . . .

Sloane on-line Encyclopedia of integer sequences:

“A006206 Number of aperiodic binary necklaces of length  $n$  with no subsequence 00, excluding the necklace “0”. (Formerly M0317)

1, 1, 1, 1, 2, 2, 4, 5, 8, 11, 18, 25, 40, 58, 90, 135, 210, 316, 492, 750, 1164, 1791, 2786, 4305, 6710, 10420, 16264, 25350, 39650, 61967, 97108, “

The partial sums (sequence  $g_n + 1$  appears there as well as A001461 in connection with some results in knot theory and field theory)

# Two-step classification

- ① Finding a presentation of the smallest member in a particular family,
- ② Description of particular cyclic extensions adding multiple edges.

## Theorem (Hu, N., Škovič, Wang, 2013...)

Let  $\mathcal{M} = (G; x, y)$  be a regular map of valency  $d$  with underlying graph  $X^{(m)}$ . Set  $A = \langle x^{d/m} \rangle$  and  $B = \langle x^{d/m}, y \rangle$ . Then:

## Theorem (Hu, N., Škovič, Wang, 2013...)

Let  $\mathcal{M} = (G; x, y)$  be a regular map of valency  $d$  with underlying graph  $X^{(m)}$ . Set  $A = \langle x^{d/m} \rangle$  and  $B = \langle x^{d/m}, y \rangle$ . Then:

- (i) The group  $A$  is a normal subgroup of  $G$ , the map  $\mathcal{M}/A = (G/A; xA, yA)$  is a regular embedding of  $X$ , and the natural projection  $\mathcal{M} \rightarrow \mathcal{M}/A$  is a map homomorphism bijective on the vertices.

## Theorem (Hu, N., Škovič, Wang, 2013...)

Let  $\mathcal{M} = (G; x, y)$  be a regular map of valency  $d$  with underlying graph  $X^{(m)}$ . Set  $A = \langle x^{d/m} \rangle$  and  $B = \langle x^{d/m}, y \rangle$ . Then:

- (i) The group  $A$  is a normal subgroup of  $G$ , the map  $\mathcal{M}/A = (G/A; xA, yA)$  is a regular embedding of  $X$ , and the natural projection  $\mathcal{M} \rightarrow \mathcal{M}/A$  is a map homomorphism bijective on the vertices.
- (ii)  $\mathcal{M}' = (B; x^{d/m}, y)$  is a dipole map isomorphic to  $\mathcal{D}(m, e)$  for some integer  $e$  such that  $e^2 \equiv 1 \pmod{m}$ .

## Theorem (Hu, N., Škovič, Wang, 2013...)

Let  $\mathcal{M} = (G; x, y)$  be a regular map of valency  $d$  with underlying graph  $X^{(m)}$ . Set  $A = \langle x^{d/m} \rangle$  and  $B = \langle x^{d/m}, y \rangle$ . Then:

- (i) The group  $A$  is a normal subgroup of  $G$ , the map  $\mathcal{M}/A = (G/A; xA, yA)$  is a regular embedding of  $X$ , and the natural projection  $\mathcal{M} \rightarrow \mathcal{M}/A$  is a map homomorphism bijective on the vertices.
- (ii)  $\mathcal{M}' = (B; x^{d/m}, y)$  is a dipole map isomorphic to  $\mathcal{D}(m, e)$  for some integer  $e$  such that  $e^2 \equiv 1 \pmod{m}$ .
- (iii)  $G = \langle x, y \mid x^{km} = y^2 = 1, yx^ky = x^{ek}, \dots \rangle$ , where  $e^2 \equiv 1 \pmod{m}$  and  $k = d/m$  is the valency of  $X$ .

## Theorem (Hu, N., Škovič, Wang, 2013...)

Let  $\mathcal{M} = (G; x, y)$  be a regular map of valency  $d$  with underlying graph  $X^{(m)}$ . Set  $A = \langle x^{d/m} \rangle$  and  $B = \langle x^{d/m}, y \rangle$ . Then:

- (i) The group  $A$  is a normal subgroup of  $G$ , the map  $\mathcal{M}/A = (G/A; xA, yA)$  is a regular embedding of  $X$ , and the natural projection  $\mathcal{M} \rightarrow \mathcal{M}/A$  is a map homomorphism bijective on the vertices.
- (ii)  $\mathcal{M}' = (B; x^{d/m}, y)$  is a dipole map isomorphic to  $\mathcal{D}(m, e)$  for some integer  $e$  such that  $e^2 \equiv 1 \pmod{m}$ .
- (iii)  $G = \langle x, y \mid x^{km} = y^2 = 1, yx^ky = x^{ek}, \dots \rangle$ , where  $e^2 \equiv 1 \pmod{m}$  and  $k = d/m$  is the valency of  $X$ .
- (iv) If  $\mathcal{M}$  is not bipartite, then  $e \equiv 1 \pmod{m}$ .

## Parameter $e$ in a nilpotent map coming from a 2-group

We have a relation  $yx^{2^t}y = x^{e2^t}$ , where  $k = 2^t$  and  $m = 2^s$ .

Since  $e^2 = 1 \pmod{m}$ , only four values  $e = \pm 1$  and  $e = 2^{s-1} \pm 1$  are admissible.

# Parameter $e$ in a nilpotent map coming from a 2-group

We have a relation  $yx^{2^t}y = x^{e2^t}$ , where  $k = 2^t$  and  $m = 2^s$ .

Since  $e^2 = 1 \pmod m$ , only four values  $e = \pm 1$  and  $e = 2^{s-1} \pm 1$  are admissible.

**Example:** Consider a family of regular embeddings of  $C_n^{(m)}$ , where both  $n, m$  are powers of two. By the classification (Hu, N., Škoviera and Wang) for each fixed  $n \geq 4$  and  $m$  there are exactly 8 such maps:

$$G(n, m; e, f) = \langle x, y \mid x^{2m} = y^2 = 1, y^{-1}x^2y = (x^2)^e, (xy)^n = (x^2)^f \rangle,$$

where  $e^2 = 1 \pmod m$  and  $f = \frac{1}{4}(e+1)n$  or  $f = \frac{1}{4}((e+1)n + 2m)$ .

# Maximal discrete actions of genus $g$

Let  $\lambda(g)$  denote the order of a largest group of conformal automorphisms of a compact Riemann surface of genus  $g$ . Accola (1968) and MacLachlan (1969) independently proved that

$$8(g + 1) \leq \lambda(g) \leq 84(g - 1)$$

for  $g \geq 2$  and there are infinitely many integers  $g \geq 2$  for which the equality  $\lambda(g) = 8(g + 1)$  holds.

# Maximal nilpotent actions of genus $g$

If we take  $n = 4$ ,  $e = -1$ , and  $f = 0$ , we get that the genus of  $\mathcal{C}(4, m; -1, 0)$  is  $g = m - 1$  with the automorphism group  $G$  of order  $|G| = 8(g + 1)$ , the lower bound of  $\lambda(g)$ .

The same problem for nilpotent groups has a solution

$$8(g + 1) \leq \lambda_{nil}(g) \leq 16(g - 1),$$

see Zamarrodian 1985, both extremal maps come from 2-groups.

# Nilpotent dessins

A *dessin* is a bipartite map  $M$  on an orientable surface with a prescribed coloring of vertices.

# Nilpotent dessins

A *dessin* is a bipartite map  $M$  on an orientable surface with a prescribed coloring of vertices.

Automorphism group  $Aut_0(M)$  of a dessin is **the colour-preserving subgroup** of  $Aut(M)$ ,

# Nilpotent dessin

A *dessin* is a bipartite map  $M$  on an orientable surface with a prescribed coloring of vertices.

Automorphism group  $Aut_0(M)$  of a dessin is **the colour-preserving subgroup** of  $Aut(M)$ ,

A dessin is *regular* if  $Aut_0(M)$  is regular on the edges.

# Nilpotent dessin

A *dessin* is a bipartite map  $M$  on an orientable surface with a prescribed coloring of vertices.

Automorphism group  $Aut_0(M)$  of a dessin is **the colour-preserving subgroup** of  $Aut(M)$ ,

A dessin is *regular* if  $Aut_0(M)$  is regular on the edges.

A dessin is abelian, nilpotent, if it is regular and  $Aut_0(M)$  is abelian, nilpotent (respectively).

# Representation of regular dessins

If  $D$  is regular, then we can label the edges of  $D$  by the elements of  $G = \text{Aut}(D)$ . Let  $\langle x \rangle$  and  $\langle y \rangle$  be generators of the stabilisers of two adjacent vertices (one is black the other is white). Hence we have an associated triple  $(G; x, y)$ , where  $G = \langle x, y \rangle$ .

# Representation of regular dessins

If  $D$  is regular, then we can label the edges of  $D$  by the elements of  $G = \text{Aut}(D)$ . Let  $\langle x \rangle$  and  $\langle y \rangle$  be generators of the stabilisers of two adjacent vertices (one is black the other is white). Hence we have an associated triple  $(G; x, y)$ , where  $G = \langle x, y \rangle$ .

Vice-versa, given  $(G; x, y)$  we set  $E = G$ , define the black vertices to be the (right) cosets of  $\langle x \rangle$  and white vertices to be the cosets of  $\langle y \rangle$ . The edge-vertex incidence is given by the containment relation.

# Nilpotent maps versus nilpotent dessins

## Theorem

*A nilpotent non-abelian map determines a nilpotent dessin.*

Vice-versa it is not true. So, nilpotent dessins form a broader family of structures.

# Decomposition theorem for nilpotent dessins

## Theorem (Decomposition Theorem)

*Let  $G \cong \prod_{i=1}^k G_i$  be the direct product decomposition of a finite two-generated nilpotent group  $G$  where  $G_i$  are the Sylow  $p_i$ -subgroups of  $G$ .*

*Then a regular dessin  $D = (G, x, y)$  is the parallel product of regular dessins  $D_i = (G_i, x_i, y_i)$  where  $x_i$  and  $y_i$  are the images of  $x$  and  $y$  under the natural projections  $\pi_i : G \rightarrow G_i$ .*

Conclusion: It is sufficient to restrict investigation to  **$p$ -dessins**.

# Shadow dessins

## Theorem

*Let  $D = (G, x, y)$  be a nilpotent regular dessin, and let  $\bar{D} = (\bar{G}, \bar{x}, \bar{y})$  be its shadow dessin. If  $c(G) = c \geq 2$ , then  $c - 1 \leq c(\bar{G}) \leq c$ . Conversely, if  $c(\bar{G}) = c$ , then  $c \leq c(G) \leq c + 1$ .*

Conclusion: Classification can be done in the two steps, first considering dessins without multiple edges.

# The beginning: Abelian dessins

## Theorem

*Up to duality swapping the black and white vertices the abelian dessins with  $\text{Aut}(D) \cong Z_{p^a} \times Z_{p^b}$ ,  $0 \leq a \leq b$  are in 1-1 correspondence with the pairs  $(c, e)$  of integers such that*

$$0 \leq c \leq b - a, \quad \text{and } e \in Z_{p^c}^*,$$

*in which case  $\text{Aut}(D)$  has presentation*

$$\langle x, y \mid x^{p^b} = y^{p^{a+c}} = [x, y] = 1, y^{p^a} = x^{ep^{b-c}} \rangle.$$

# How many abelian dessins there are?

## Theorem

*Up to isomorphism, the number of regular dessins  $A$  with  $\text{Aut}(A) \cong Z_n \oplus Z_m$  where  $0 \leq n \leq m$  is equal to  $\psi(m/n)$  where  $\psi$  is the Dedekind's totient function.*

# How many abelian dessins there are?

## Theorem

*Up to isomorphism, the number of regular dessins  $A$  with  $\text{Aut}(A) \cong Z_n \oplus Z_m$  where  $0 \leq n \leq m$  is equal to  $\psi(m/n)$  where  $\psi$  is the Dedekind's totient function.*

**Remark.** The Dedekind totient function  $\psi(m) = m \prod_{p|m} (1 + \frac{1}{p})$  was introduced by Dedekind in connection with his investigation of modular forms. It appears in many other contexts, for instance in the formula describing the generating function of the Riemann zeta function.

# Properties of abelian dessins

- 1 All are reflexible,

# Properties of abelian dessins

- ① All are reflexible,
- ② the type is  $\{2mn/d, mn\}$ ,

# Properties of abelian dessins

- 1 All are reflexible,
- 2 the type is  $\{2mn/d, mn\}$ ,
- 3 the underlying graph is  $K_{n,n}^{(m)}$ ,

# Properties of abelian dessins

- 1 All are reflexible,
- 2 the type is  $\{2mn/d, mn\}$ ,
- 3 the underlying graph is  $K_{n,n}^{(m)}$ ,
- 4 the shadow map is the standard embedding of  $K_{n,n}$  (Hidalgo (2013))

## Symmetric dessins of class 2

### Theorem

*Simple symmetric  $p$ -dessins of class two are in correspondence with quadruples  $(a, b, c, u)$  such that*

$$\max(0, 1 + c - a) \leq b \leq c \leq a \leq 2c - b, u \in \mathbb{Z}_{p^{a-c}}^*$$

*in which case it has presentation*

$$\langle x, y \mid x^{p^a} = y^{p^a} = [x, z] = [y, z] = 1, z^{up^b} = x^{-p^c} u^{p^c}, z = [x, y] \rangle$$

# Enumeration of simple dessins of class 2

## Theorem

*The number of symmetric simple dessins is given by the function*

$$F(p, a) = (p + 1)(p^{a/2} - 1)/(p - 1),$$

*if  $a$  is even, and*

$$F(p, a) = (2p^{\frac{a+1}{2}} - p - 1)/(p - 1),$$

*if  $a$  is odd.*

# Further results

- ① complete classification of symmetric dessins of class 2,
- ② regular embeddings of  $K_{n,n}^{(m)}$  over standard embedding of  $K_{n,n}$ ,
- ③ regular embeddings of  $K_{p^e,p^e}^{(m)}$ , for a prime  $p > 2$ ,
- ④ regular embeddings of  $K_{2^e,2^e}^{(m)}$  incomplete.

# Open problems

- ① Labute proved a more general results on the size of  $\Gamma_n/\Gamma_{n+1}$  for the one-relator group  $\langle x, y | y^p = 1 \rangle$ . We have employed it just in case  $p = 2$ . Is there a room for generalisation?
- ② It seems regular dessins with the underlying graphs  $K_{n,n}^{(m)}$  play important role. Can we complete the classification?
- ③ A project: Groups or graphs giving rise to a unique regular dessin (map).

# A sample of the uniqueness phenomenon

An integer  $n$  is called singular, if  $\gcd(n, \varphi(n)) = 1$ .

## Theorem

*The graph  $K_{n,n}^{(m)}$  has a unique regular embedding iff  $m = 1$  and  $n$  is singular, or  $m = 2$  and  $n$  is odd and singular.*

This generalises an older statement on the uniqueness of regular embeddings of  $K_{n,n}$  by Jones, N., Škoviera.