

Characteristic polynomial of the transfer matrix:

$$\begin{aligned}
p(x) = & x^{18}(x-1)^2(x+1)(x^{71}-x^{69}-2x^{68}-x^{67}+2x^{66}+2x^{65}+x^{64} \\
& -x^{63}-x^{62}-x^{61}-x^{60}-x^{59}+2x^{58}+5x^{57}+3x^{56}-2x^{55}-10x^{54} \\
& -3x^{53}-2x^{52}+6x^{51}+6x^{50}+x^{49}+9x^{48}-3x^{47}-7x^{46}-8x^{45} \\
& -8x^{44}+10x^{43}+6x^{42}+8x^{41}-5x^{40}-12x^{39}+7x^{38}-7x^{37}+7x^{36} \\
& +x^{35}-3x^{34}+10x^{33}+x^{32}-6x^{31}-2x^{30}-10x^{29}-3x^{28}+2x^{27} \\
& +9x^{26}-3x^{25}+14x^{24}-8x^{23}-7x^{21}+9x^{20}+3x^{19}-4x^{18}-10x^{17} \\
& -7x^{16}+12x^{15}+7x^{14}+2x^{13}-12x^{12}-4x^{11}-2x^{10}+5x^9+x^7-7x^6 \\
& +7x^5-4x^4+12x^3-6x^2+3x-6)
\end{aligned}$$

Main irreducible part thereof:

$$\begin{aligned}
q(x) = & x^{71}-x^{69}-2x^{68}-x^{67}+2x^{66}+2x^{65}+x^{64}-x^{63}-x^{62}-x^{61} \\
& -x^{60}-x^{59}+2x^{58}+5x^{57}+3x^{56}-2x^{55}-10x^{54}-3x^{53}-2x^{52} \\
& +6x^{51}+6x^{50}+x^{49}+9x^{48}-3x^{47}-7x^{46}-8x^{45}-8x^{44}+10x^{43} \\
& +6x^{42}+8x^{41}-5x^{40}-12x^{39}+7x^{38}-7x^{37}+7x^{36}+x^{35}-3x^{34} \\
& +10x^{33}+x^{32}-6x^{31}-2x^{30}-10x^{29}-3x^{28}+2x^{27}+9x^{26}-3x^{25} \\
& +14x^{24}-8x^{23}-7x^{21}+9x^{20}+3x^{19}-4x^{18}-10x^{17}-7x^{16}+12x^{15} \\
& +7x^{14}+2x^{13}-12x^{12}-4x^{11}-2x^{10}+5x^9+x^7-7x^6+7x^5-4x^4+12x^3 \\
& -6x^2+3x-6
\end{aligned}$$

Biggest by absolute value root of $q(x)$ (and also of $p(x)$):

$$\lambda = 1.30357726903429639125709911215255189073070250465940487575486139\dots$$

All complex roots of $q(x)$:

