

ON THE STRUCTURE OF THE STABLE NORM OF PERIODIC METRICS

D. BURAGO, S. IVANOV, AND B. KLEINER

ABSTRACT. We study the differentiability of the stable norm $\|\cdot\|$ associated with a \mathbf{Z}^n -periodic metric on \mathbf{R}^n . Extending one of the main results of [Ba2], we prove that if $p \in \mathbf{R}^n$ and the coordinates of p are linearly independent over \mathbf{Q} , then there is a linear 2-plane V containing p such that the restriction of $\|\cdot\|$ to V is differentiable at p . We construct examples where $\|\cdot\|$ is not differentiable at a point with coordinates linearly independent over \mathbf{Q} .

INTRODUCTION

0.1. In this paper we study the large-scale geometry of an abelian cover of a compact Riemannian manifold M^m . If \hat{M} is a normal cover of M with deck group \mathbf{Z}^n , then \hat{M} is within finite Gromov-Hausdorff distance from an n -dimensional normed vector space, [Bu1], which we denote $(\mathbf{R}^n, \|\cdot\|)$. We call this the *stable norm*.

We denote by B and F the unit ball and the unit sphere of the norm $\|\cdot\|$, i.e. $B = \{v \in \mathbf{R}^n : \|v\| \leq 1\}$, $F = \{v \in \mathbf{R}^n : \|v\| = 1\}$, respectively. Our subject is the local structure of the surface F , namely, the shape of its tangent cone at certain points. This is a part of the general (and wide open) question: what norms can arise as stable norms of periodic Riemannian metrics?

In [Ba2] V. Bangert – inspired by [AuMa] – made substantial progress on this question in the case of two dimensional tori (analogous results were obtained independently by F. Nazarov [Na]). [Ba2] proved that the stable norm is differentiable at every irrational point¹ in \mathbf{R}^2 , and that it is differentiable at a rational point only if the torus is foliated by closed geodesics representing the corresponding element of $\mathbf{Z}^2 \simeq H_1(T^2; \mathbf{Z})$.

In this paper, we consider the higher dimensional case, and the regularity of the stable norm in irrational directions. The following theorem is a partial generalization of V. Bangert's result [Ba2], see also [Ba4].

Theorem 1. *Assume that $m \leq n$, and p is an irrational point in \mathbf{R}^n . Then the stable norm $\|\cdot\|$ is differentiable in at least one nonradial direction at p ; in other*

The first author was supported by NSF grant DMS-95-05175. The second author was supported by RFFR grant 96-01-00674 and CRDF grant RM1-169. The third author was supported by NSF grants DMS-95-05175 and DMS-9626911.

¹A point $p \in \mathbf{R}^n$ is irrational if its coordinates are linearly independent over \mathbf{Q} .

words, the tangent cone to F at $\frac{p}{\|p\|}$ splits as a metric product of \mathbf{R} with another cone.

The examples described in the next theorem show that the stable norm can be nondifferentiable at irrational points; hence the most optimistic attempt to generalize [Ba2] fails.

Theorem 2. *For every k , there is an n such that for almost every irrational point $p \in \mathbf{R}^n$, there is a C^k -smooth Riemannian metric on n -torus whose stable norm is not differentiable at p .*

It turns out that the (non)differentiability of the asymptotic norm is directly related to the structure of certain minimizing geodesics in \hat{M} and their interplay with a class of distance-like functions. Differentiability – or more precisely directional differentiability – correlates with the existence of curves which deviate arbitrarily far from a minimizing geodesic with bounded additional “cost”. Such curves must lie in a small neighborhood of the family of minimizers associated with the point of \mathbf{R}^n ; the idea behind theorem 1 was that such curves may be constructed by joining long pieces of minimizers with short segments. We note that even for smooth metrics, the set of minimizing geodesics can be badly behaved. In the example of theorem 2, the surface formed by the set of minimizing geodesics is highly corrugated; every curve that stays close to it must be “long”.

Theorems 1 and 2 give some new information about possible singularities of the stable norm, but the picture is still far from complete. The following two questions remain open:

1. Is the finite smoothness essential for the examples in theorem 2, or are there similar C^∞ examples?
2. If the stable norm is smooth and uniformly convex on an open set, is \hat{M} foliated by minimizing geodesics?

To simplify notation, we will deal with the case where M is diffeomorphic to T^n , and \hat{M} is the universal cover of M ; therefore we may identify \hat{M} with \mathbf{R}^n . Although this identification is non-invariant, the image of $\mathbf{Z}^n \subset \mathbf{R}^n$ under this identification is. Then the stable norm $\|\cdot\|$ is given by

$$\|v\| = \lim_{\lambda \rightarrow \infty} \frac{\rho(0, \lambda v)}{\lambda}$$

where $\rho(\cdot, \cdot)$ is the distance function on $\hat{M} \simeq \mathbf{R}^n$. Sometimes this norm is also called *limit norm* or *asymptotic norm* of a metric ρ . The Banach space $(\mathbf{R}^n, \|\cdot\|)$ approximates the metric space (\mathbf{R}^n, ρ) in a very strong sense: there exists a constant $C = C(\rho)$ such that

$$(1) \quad \forall x, y \in \mathbf{R}^n \quad \left| \|x - y\| - \rho(x, y) \right| \leq C$$

(see [Bu1]). This estimate refines the statement that the Gromov–Hausdorff distance between (\mathbf{R}^n, ρ) and $(\mathbf{R}^n, \|\cdot\|)$ is finite.

We will use ρ to denote the periodic metric under consideration and $\bar{\rho}$ for the corresponding metric on the torus $T^n = \mathbf{R}^n/\mathbf{Z}^n$. We denote by UTT^n and $UT\mathbf{R}^n$ the unit tangent bundles for metrics $\bar{\rho}$ and ρ . All geodesics are parameterized by arc length. We call a geodesic γ in $(T^n, \bar{\rho})$ minimal if its lift $\tilde{\gamma}$ is minimal, i.e. if $\rho(\tilde{\gamma}(a), \tilde{\gamma}(b)) = |a - b|$ for all $a, b \in \mathbf{R}$.

ACKNOWLEDGEMENTS

The first author would like to thank V. Bangert and F. Nazarov for discussions which to some extent formed his vision of the problem. He also would like to thank A. Katok and N. Simanai for their patience while listening to the final version of the argument.

1. ROTATION VECTORS AND UNIFORMLY RECURRENT GEODESICS

1.1. We define the *direction at infinity* $D(\gamma) \in \mathbf{R}^n$ for a geodesic $\gamma: \mathbf{R} \rightarrow (\mathbf{R}^n, \rho)$ by

$$D(\gamma) = \lim_{t \rightarrow \infty} \frac{\gamma(t) - \gamma(0)}{t}$$

if the limit exists. Since $\rho(\gamma(t), \gamma(0)) \leq t$ for all $T \in \mathbf{R}$, we have

$$\|D(\gamma)\| = \lim_{t \rightarrow \infty} \frac{\|\gamma(t) - \gamma(0)\|}{t} = \lim_{t \rightarrow \infty} \frac{\rho(\gamma(t), \gamma(0))}{t} \leq 1$$

(the second equality follows from (1)). The *rotation vector* (see [?]) $R(\gamma) \in \mathbf{R}^n$ of a geodesic $\gamma: \mathbf{R} \rightarrow (T^n, \bar{\rho})$ is defined by $R(\gamma) = D(\tilde{\gamma})$ where $\tilde{\gamma}$ is a lift of γ .

Recall that a geodesic $\gamma: \mathbf{R} \rightarrow (T^n, \bar{\rho})$ is *uniformly recurrent* if for any $t_0 \in \mathbf{R}$ the trajectory $\{\gamma'(t) : t \in \mathbf{R}\}$ visits any neighborhood of $\gamma'(t_0)$ in UTT^n with a positive frequency. The purpose of this section is to prove the following

1.2. Proposition. *If v is an extreme point of B (i.e. v is not the midpoint of a line segment contained in B), then there is a uniformly recurrent geodesic $\gamma: \mathbf{R} \rightarrow (T^n, \bar{\rho})$ with $R(\gamma) = v$.*

1.3. Remark. It follows from 2.4 that a geodesic γ given by Proposition 1.2 is a minimal one. Thus we obtain a kind of existence statement for minimal geodesics with a given rotation vector. In general, not every vector $v \in F$ can be obtained as a rotation vector of a minimal geodesic. For $n \geq 3$ there exist examples of periodic metrics for which B is a polyhedron and its vertices are only possible rotation vectors of minimal geodesics (see [Ba1]).

1.4. We will prove Proposition 1.2 using the technique of minimal measures introduced by J. Mather [M2]. In fact, we only adopt the basic constructions of [M2] to our settings.

We may also view the rotation vector of a geodesic as a function of its velocity vector, i.e. for $w \in UTT^n$ we set $R(w) = R(\gamma)$ where γ is a geodesic in $(T^n, \bar{\rho})$ with $\gamma'(0) = w$. This way R becomes a function defined on a subset of UTT^n . Define a map $\omega: TT^n \rightarrow \mathbf{R}^n$ as the second projection of the natural decomposition $TT^n \cong T^n \times \mathbf{R}^n$. The rotation vector of a geodesic γ in $(T^n, \bar{\rho})$ may be written in the form

$$(2) \quad R(\gamma) = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \tilde{\gamma}' = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \omega \circ (\gamma').$$

(here $\tilde{\gamma}$ denotes a lift of γ in \mathbf{R}^n). Thus the function R is the average along trajectories of the function ω .

Let m be a finite Borel measure on UTT^n . We define its rotation vector $\mathcal{R}(m) \in \mathbf{R}^n$ by the formula

$$\mathcal{R}(m) = \int_{UTT^n} \omega \, dm.$$

Clearly $\mathcal{R}(m)$ is a linear function of m . We call a measure m on UTT^n *invariant* if it is preserved by the geodesic flow of $(T^n, \bar{\rho})$. If m is an invariant probability measure then (by (2) and the Birkhoff ergodic theorem) the function R is defined m -almost everywhere, and

$$\mathcal{R}(m) = \int_{UTT^n} R(w) dm(w).$$

In particular, $\|\mathcal{R}(m)\| \leq 1$. The first observation about invariant measures is the following

1.3. Lemma. *For every $v \in F$ there is an invariant probability measure m with $\mathcal{R}(m) = v$.*

Proof. Let (λ_i) be a sequence of positive numbers, $\lambda_i \rightarrow \infty$. For each i let $\gamma_i: [0, l_i] \rightarrow (T^n, \bar{\rho})$ be a minimal geodesic whose lift joins points 0 and $\lambda_i v$ in (\mathbf{R}^n, ρ) . Here $l_i = \rho(0, \lambda_i v)$. Then consider a probability measure uniformly distributed along the segment $[0, l_i] \subset \mathbf{R}$ and let m_i be the image of that measure in UTT^n under the map $t \mapsto \gamma_i'(t)$. Some subsequence of (m_i) converges weakly in the space of probability measures on UTT^n . We may assume that the original sequence (m_i) converges to some measure m . It is trivial that m is an invariant measure. Then

$$\mathcal{R}(m) = \int_{UTT^n} \omega dm = \lim \int_{UTT^n} \omega dm_i = \lim \frac{\lambda_i v}{l_i} = \frac{v}{\|v\|}$$

(the last equality follows from the definition of the stable norm). Since $\|v\| = 1$, the lemma follows. \square

1.4. Now suppose that v is an extreme point of B (i.e. no line segment contained in B has its midpoint at v). Let $\mathcal{M}(v)$ denote the set of all invariant probability measures m with $\mathcal{R}(m) = v$. This set is convex and compact (with respect to the weak topology). By the Krein–Milman theorem there is a measure m which is an extreme point of $\mathcal{M}(v)$. Such a measure m is ergodic with respect to the geodesic flow. Indeed, if $0 < \lambda < 1$ and $m = \lambda m_1 + (1 - \lambda)m_2$ for some invariant probability measures m_1 and m_2 , then $v = \mathcal{R}(m) = \lambda \mathcal{R}(m_1) + (1 - \lambda)\mathcal{R}(m_2)$, so the extremeness of v in B implies that $\mathcal{R}(m_1) = \mathcal{R}(m_2) = v$. Since m is an extreme point of $\mathcal{M}(v)$ it follows that $m_1 = m_2$.

Let U be an open subset of UTT^n with $m(U) > 0$. The ergodicity of m implies that m -almost every trajectory of the geodesic flow visits U with positive frequency. Since the topology of UTT^n has a countable base, m -almost every trajectory does this for all open sets of positive measure. Note that m -almost every trajectory is contained in $\text{supp}(m)$, the support of measure m . Since for an open $U \subset UTT^n$ the condition $U \cap \text{supp}(m) \neq \emptyset$ implies $m(U) > 0$, it follows that m -almost every trajectory is uniformly recurrent. On the other hand, $R(w) = \mathcal{R}(m) = v$ for m -almost all $w \in UTT^n$ since m is ergodic. This completes the proof of Proposition 1.2.

2. GENERALIZED COORDINATES AND MINIMIZERS

2.1. If $L: \mathbf{R}^n \rightarrow \mathbf{R}$ is a linear function we let $\|L\|$ denote its norm in the space $(\mathbf{R}^n, \|\cdot\|)^*$, i.e. $\|L\| = \max\{|L(x)| : \|x\| = 1\}$. We say that a linear function L *supports* B at a point $v \in F$ if $\|L\| = 1$ and $L(v) = 1$. Geometrically it means that

a hyperplane $L^{-1}(1)$ touches the surface F at p . We denote by $F^*(v)$ the set of linear functions supporting B at v , and by F^* the set of all linear functions L with $\|L\| = 1$.

Let $L \in F^*$ and let f be a real-valued function on (\mathbf{R}^n, ρ) . We say that f is a *generalized coordinate* associated with L if

(2.1.1) f is a 1-Lipschitz function with respect to ρ , i.e. $|f(x) - f(y)| \leq \rho(x, y)$ for all $x, y \in \mathbf{R}^n$.

(2.1.2) The function $(f - L)$ is \mathbf{Z}^n -periodic, i.e. for every $x \in \mathbf{R}^n$ and $h \in \mathbf{Z}^n$ we have $f(x + h) = f(x) + L(h)$.

Such function were used in [BuI2] to make volume estimates. We extract a construction from [BuI2] in the following statement.

2.2. Proposition. *For any $L \in F^*$ there is a generalized coordinate associated with L .*

Proof. Define a function f on \mathbf{R}^n by

$$f(x) = \limsup_{\|y\| \rightarrow \infty} (L(y) - \rho(x, y)).$$

First, we need to prove that all values of f are finite. This follows from (1) and the equality

$$\limsup_{\|y\| \rightarrow \infty} (L(y) - \|x - y\|) = L(x)$$

which holds since $\|L\| = 1$. Then f is 1-Lipschitz as it is the supremum of a family of 1-Lipschitz functions. Finally, for any $x \in \mathbf{R}^n$ and $h \in \mathbf{Z}^n$ we have

$$\begin{aligned} f(x + h) &= \limsup_{\|y\| \rightarrow \infty} (L(y) - \rho(x + h, y)) \\ &= \limsup_{\|y-h\| \rightarrow \infty} (L(y - h) - \rho(x, y - h)) + L(h) = f(x) + L(h), \end{aligned}$$

so f satisfies (2.1.2). \square

2.3. Let f be a generalized coordinate and let $\gamma: \mathbf{R} \rightarrow (\mathbf{R}^n, \rho)$ be a geodesic. We say that γ is *calibrated by f* , or that γ is an *f -calibrated* [HaLa], if

$$(3) \quad f(\gamma(b)) - f(\gamma(a)) = b - a$$

for all $a, b \in \mathbf{R}$.

Note that for an arbitrary unit-speed curve γ in (\mathbf{R}^n, ρ) we have

$$f(\gamma(b)) - f(\gamma(a)) \leq \rho(\gamma(a), \gamma(b)) \leq |b - a|$$

for all $a, b \in \mathbf{R}$. Therefore if γ is f -calibrated then γ is a minimal geodesic. It is clear that being f -calibrated is actually a local property: if γ is not f -calibrated then for any $\varepsilon > 0$ there exists $a \in \mathbf{R}$ such that (3) fails for $[a, b] = [a, a + \varepsilon]$. From (2.1.2) it follows that any integer translate of an f -calibrated geodesic is also an f -calibrated.

2.4. Proposition. *Let $\gamma: \mathbf{R} \rightarrow (\mathbf{R}^n, \rho)$ be a lift of a uniformly recurrent geodesic with $D(\gamma) = v \in F$, and $L \in F^*(v)$. Then γ is calibrated by any generalized coordinate associated with L .*

Proof. Let f be a generalized coordinate associated with L , and suppose that γ is not f -calibrated. We may assume that (3) fails for $a = 0$ and $b = 1$, say $f(\gamma(1)) - f(\gamma(0)) = 1 - \varepsilon$ where $\varepsilon > 0$. Let U be a neighborhood of $\gamma'(0)$ in $UT\mathbf{R}^n$ such that for every geodesic γ_1 in (\mathbf{R}^n, ρ) with $\gamma_1'(0) \in U$ we have $\rho(\gamma_1(t), \gamma(t)) < \varepsilon/3$ for all $t \in [0, 1]$. Since γ is a lift of a uniformly recurrent geodesic, there exists a sequence $(t_i)_{i=1}^\infty$ of real numbers such that for all i we have $t_{i+1} \geq t_i + 1$, $t_i/i \leq T_0 < \infty$, and $\gamma'(t_i) \in U + h_i$ for some $h_i \in \mathbf{Z}^n$. By (2.1.2) and the choice of U we have

$$f(\gamma(t_i + 1)) - f(\gamma(t_i)) \leq 1 - \varepsilon/3$$

for each i . Summing up these inequalities for $i = 1, \dots, m$, together with obvious ones

$$f(\gamma(t_{i+1})) - f(\gamma(t_i)) \leq t_{i+1} - t_i$$

we obtain that

$$f(\gamma(t_m + 1)) - f(\gamma(0)) \leq t_m + 1 - \frac{i\varepsilon}{3} \leq \left(1 - \frac{\varepsilon}{3T_0}\right) t_m + 1.$$

Therefore

$$\limsup_{t \rightarrow \infty} \frac{f(\gamma(t)) - f(\gamma(0))}{t} \leq 1 - \frac{\varepsilon}{3T_0} < 1.$$

Since $|f - L|$ is bounded, one may replace f by L in this estimate. On the other hand,

$$\lim_{t \rightarrow \infty} \frac{L(\gamma(t)) - L(\gamma(0))}{t} = L(D(\gamma)) = L(v) = 1$$

(the first equality follows from the definition of $D(\gamma)$, see 1.1). This contradiction proves the proposition. \square

We will apply Proposition 2.4 at a point $v \in F$ where B has several different linear support functions (this may happen if F is not smooth at v), so the same geodesic γ will be calibrated by the generalized coordinates associated with each of these functions. The following Proposition 2.5 tells us that in this case all the generalized coordinates calibrating γ have similar local behavior near γ .

2.5 Proposition. *Let ρ be a C^3 metric. Then there is a constant $C = C(\rho)$ with the following property. If a geodesic γ is calibrated by two generalized coordinates f_1 and f_2 (possibly associated with different linear functions), then for $d = f_1(\gamma(0)) - f_2(\gamma(0))$ the inequality*

$$|f_1(x) - f_2(x) - d| \leq C \cdot \rho(x, \gamma)^2$$

holds for all $x \in \mathbf{R}^n$. Here $\rho(x, \gamma)$ denotes $\inf\{\rho(x, \gamma(t)) : t \in \mathbf{R}\}$.

Proof. It suffices to prove the statement only in the case $d = 0$ (just add a constant to f_2). Pick constants $\varepsilon > 0$ and $r > \varepsilon$ such that $r + \varepsilon$ is less than the injectivity

radius of (\mathbf{R}^n, ρ) . Fix $t \in \mathbf{R}$ and denote $c = f_1(\gamma(t)) = f_2(\gamma(t))$. Since f_1 and f_2 are 1-Lipschitz functions, we have

$$f_i(\gamma(t+r)) - \rho(\gamma(t+r), x) \leq f_i(x) \leq f_i(\gamma(t-r)) + \rho(\gamma(t-r), x),$$

or

$$f_{\text{low}}(x) := c + r - \rho(\gamma(t+r), x) \leq f_i(x) \leq c - r + \rho(\gamma(t-r), x) := f_{\text{up}}(x)$$

for any $x \in \mathbf{R}^n$, $i = 1, 2$. Both f_{low} and f_{up} are C^2 -smooth functions of x within ε -neighborhood of $\gamma(t)$, and their values and first derivatives at $\gamma(t)$ coincide. (The values are equal to c and the gradients are equal to $\gamma'(t)$). So if $C > 2/\varepsilon$ and C is an upper bound for the second derivative of a function $\rho(x_0, \cdot)$ between its values $r - \varepsilon$ and $r + \varepsilon$, then

$$|f_1(x) - f_2(x)| \leq f_{\text{up}}(x) - f_{\text{low}}(x) \leq C \cdot \rho(x, \gamma(t))^2$$

Since t is arbitrary, the proposition follows. \square

3. A TANGENT CONE OF B AT AN IRRATIONAL POINT

3.1. We call a vector $v \in \mathbf{R}^n$ *irrational* if its coordinates are linearly independent over \mathbf{Q} : v is not an irrational vector if and only if there is a nonzero linear function $L \in (\mathbf{Q}^n)^* \subset (\mathbf{R}^n)^*$ such that $L(v) = 0$. The latter description implies that non-irrational vectors form a set of zero measure in \mathbf{R}^n since this set is a union of countably many hyperplanes. If v is an irrational vector then λv is irrational for any $\lambda \neq 0$.

For $v \in \mathbf{R}^n$ we denote by $\langle v \rangle$ the line $\{\lambda v : \lambda \in \mathbf{R}\} \subset \mathbf{R}^n$. If v is an irrational vector the union of lines $\langle v \rangle + \mathbf{Z}^n := \bigcup_{h \in \mathbf{Z}^n} (\langle v \rangle + h)$ is dense in \mathbf{R}^n .

3.2. For $v \in F$ we denote by $C_v(B)$ the tangent cone of B at v . We define a tangent cone as an intersection of half-spaces

$$(4) \quad C_v(B) = \bigcap_{L \in F^*(v)} \{x \in \mathbf{R}^n : L(x) \leq 0\}$$

A convex cone is called *sharp* if it contain no straight lines. From (4) it follows that $C_v(B)$ is sharp if and only if $F(v)$ contains n linearly independent linear functions. If $C_v(B)$ is sharp then obviously v is an extreme point of B .

3.3. Theorem. *Let ρ be a C^3 -smooth periodic Riemannian metric on \mathbf{R}^n , let B be the unit ball of its stable norm, and let v be an irrational vector in the boundary of B . Then the tangent cone $C_v(B)$ is not sharp.*

Proof. Suppose that $C_v(B)$ is a sharp cone. By Proposition 1.5 there exists a geodesic $\gamma: \mathbf{R} \rightarrow (\mathbf{R}^n, \rho)$ with $D(\gamma) = v$ which is a lift of a uniformly recurrent geodesic. Pick n linearly independent functions L_1, \dots, L_n from $F^*(v)$. For each $i = 1, \dots, n$ construct a generalized coordinate f_i associated with L_i . We may assume that $f_i(\gamma(0)) = 0$ and so $f_i(\gamma(t)) = t$ for all $t \in \mathbf{R}$. Define a map $\bar{f}: (\mathbf{R}^n, \rho) \rightarrow \mathbf{R}^{n-1}$ by

$$\bar{f} = (f_1 - f_2, f_2 - f_3, \dots, f_{n-1} - f_n).$$

For any $x \in \mathbf{R}^n$ and $h \in \mathbf{Z}^n$ we have

$$\bar{f}(x+h) = \bar{f}(x) + \bar{L}(h)$$

where the linear map $\bar{L}: \mathbf{R}^n \rightarrow \mathbf{R}^{n-1}$ is defined by

$$\bar{L} = (L_1 - L_2, L_2 - L_3, \dots, L_{n-1} - L_n).$$

Therefore for any $h \in \mathbf{Z}^n$ the values of \bar{f} are equal to $\bar{L}(h)$ along the geodesic $\gamma+h$, so we have the estimate

$$|\bar{f}(x) - \bar{L}(h)| \leq C \cdot \rho(x, \gamma+h)^2.$$

(see 2.4 and 2.5). Applying this for points x of another geodesic $\gamma+h_1$ we obtain that

$$|\bar{L}(h_1) - \bar{L}(h)| \leq C \cdot \rho(\gamma+h_1, \gamma+h)^2$$

for all $h, h_1 \in \mathbf{Z}^n$. (For two geodesics γ_1 and γ_2 we denote by $\rho(\gamma_1, \gamma_2)$ the distance between them as subsets of (\mathbf{R}^n, ρ)). Therefore

$$(5) \quad \rho(\gamma+h_1, \gamma+h) \geq \sqrt{|\bar{L}(h_1) - \bar{L}(h)|/C}$$

for all $h, h_1 \in \mathbf{Z}^n$.

The functions $f_i - L_i$ are bounded and $f_i(g(t)) = t$ for $t \in \mathbf{R}$, so $|L_i(\gamma(t)) - t| \leq C$ for some constant C not depending on t . Hence there is an $R > 0$ such that $|\gamma(t) - vt| \leq R$ for all $t \in \mathbf{R}$ (this follows from the fact that vt is the only point of \mathbf{R}^n at which the values of functions L_i are all equal to t). Thus any straight line of the form $\langle v \rangle + h$ is contained within R -neighborhood of the corresponding geodesic $\gamma+h$.

Note that the map \bar{L} is surjective and $\text{Ker } \bar{L} = \langle v \rangle$. Let U be a bounded neighborhood of a unit cube in \mathbf{R}^{n-1} , U_1 be a bounded set in \mathbf{R}^n such that $U \subset L(U_1)$, and U_2 be the R -neighborhood of U_1 . Since $\langle v \rangle + \mathbf{Z}^n$ is dense in \mathbf{R}^n and $\bar{L}(v) = 0$, the set $\bar{L}(\mathbf{Z}^n)$ is dense in \mathbf{R}^{n-1} . For a small $\varepsilon > 0$ one can find a collection of points $y_1, \dots, y_N \in U \cap \bar{L}(\mathbf{Z}^n)$ such that $N \geq (1/\varepsilon)^{n-1}$ and $|y_i - y_j| > \varepsilon$ for $i \neq j$. Let $y_i = \bar{L}(h_i)$ where $h_i \in \mathbf{Z}^n$. Consider the geodesics $\gamma+h_i$ in (\mathbf{R}^n, ρ) . They all cross the region U_2 . If U_3 is the 1-neighborhood of U_2 in (\mathbf{R}^n, ρ) , then the intersection of each of our geodesics $\gamma+h_i$ with U_3 will contain an interval of length 2.

By (5), the pairwise distances between the geodesics $\gamma+h_i$ are not less than $\sqrt{\varepsilon/C}$. Hence their tubular neighborhoods of radius $\sqrt{\varepsilon/4C}$ are disjoint. Summing up Riemannian volumes of these tubular neighborhoods we obtain a lower bound for the volume of U_3 :

$$\text{Vol}(U_3) \geq (1/\varepsilon)^n \cdot (\varepsilon/4C)^{(n-1)/2} = c_1 \varepsilon^{-(n+1)/2}$$

for some $c_1 > 0$. Since ε is arbitrarily small, it follows that $\text{Vol}(U_3) = \infty$. This is impossible since U_3 is bounded. \square

4. AN EXAMPLE

In this section we construct the examples described in theorem 2. The idea of the construction is to begin by producing a set of minimizers. This set will be (the closure of) a highly corrugated surface. The estimate (5), together with elementary geometric considerations, suggests that the cross-section of this surface may be regarded as a curve which stretches distance in the following very strong sense: the distance between the images is at least the square root of the distance between pre-images modulo 1. On the other hand, this curve has to have irrational rotation vector, and it has to be invariant under a group of diffeomorphisms. We construct this curve as a sum of two curves. One of them is a small periodic curve (with rotation vector zero), which stretches distance and persists under a group diffeomorphisms having dense orbits on the curve. The other is a smooth curve with irrational rotation vector which lies in a submanifold of high codimension. The first curve will be constructed using trigonometric series, where the following approximation condition is a technical requirement used to estimate the denominators in the coefficients of the series.

We call a number $\alpha \in \mathbf{R}$ *approximable* if for any $\varepsilon > 0$ there exists a sequence of rational numbers $\{p_k/q_k\}_{k=1}^\infty$ such that

$$|\alpha - p_k/q_k| < 1/q_k^2$$

and

$$(6) \quad q_k^{1+\varepsilon/2} < q_{k+1} < \frac{1}{2}q_k^{1+\varepsilon}$$

for all $k \geq 1$.

4.1. Lemma. *Almost all real numbers are approximable.*

Proof. For a given $\alpha \in \mathbf{R}$, every convergent p_k/q_k of the corresponding continued fraction satisfies $|\alpha - p_k/q_k| < 1/q_k^2$. It is known [?] that for almost every α the denominators of these convergents grow exponentially: there exists a positive limit $\lim_{k \rightarrow \infty} (\log q_k)/k$. For such α , it is certainly possible to find a subsequence of $\{q_k\}$ to satisfy (6). \square

Let $\gamma: \mathbf{R} \rightarrow \mathbf{R}^n$ be a continuous curve and $r > 1$ be an integer. We say that γ is *r-stretching* if there exists $\delta > 0$ such that $|\gamma(x) - \gamma(y)| \geq |x - y|^{1/r}$ whenever $|x - y| \leq \delta$.

4.2. Proposition. *Let $r > 1$ be an integer, and let $\alpha \in \mathbf{R}$ be approximable. Then there exists an r-stretching 1-periodic curve $\gamma: \mathbf{R} \rightarrow \mathbf{R}^{8r}$ such that the function*

$$x \mapsto \gamma(x + \alpha) - \gamma(x)$$

is C^1 -smooth.

Proof. Let $\varepsilon = 1/2(r - 1)$ and pick a sequence $\{p_k/q_k\}$ as in 4.1. We assume that $q_1 > 10^{4r(r-1)}$ which, by (6), implies that $q_{k+1} > 10^r q_k$ for all k .

We identify \mathbf{R}^{8r} with \mathbf{C}^{4r} and use the notation $E(t) = \exp(2\pi it) \in \mathbf{C}$ for $t \in \mathbf{R}$. Let v_k be the k^{th} basis vector of \mathbf{C}^{4r} for $1 \leq k \leq 4r$, and $v_{k+4r} = v_k$ for all $k \in \mathbf{Z}$. Define $\gamma: \mathbf{R} \rightarrow \mathbf{C}^{4n}$ by

$$\gamma(x) = \sum_{k=1}^{\infty} q_k^{-1/2r} E(q_k x) v_k.$$

Then

$$\gamma(x + \alpha) - \gamma(x) = \sum_{k=1}^{\infty} q_k^{-1/2r} E(q_k \alpha) E(q_k x) v_k .$$

The formal derivative of these series has the form

$$\sum_{k=1}^{\infty} q_k^{1-1/2r} E(q_k \alpha) E(q_k x) v_k .$$

Since $q_k \alpha - p_k < 1/q_k$, we have $E(q_k \alpha) < 2\pi q_k^{-1}$, so this formal derivative converges absolutely. It follows that the function $x \mapsto \gamma(x + \alpha) - \gamma(x)$ is C^1 .

We will prove that $|\gamma(x) - \gamma(y)| \geq |x - y|^{1/r}$ whenever $|x - y| < 1/2q_1$. Assume that $x < y < x + 1/2q_1$ and denote $\delta = y - x$. We have

$$\gamma(y) - \gamma(x) = \sum_{k=1}^{\infty} q_k^{-1/2r} E(q_k x) (E(q_k \delta) - 1) v_k .$$

The right inequality in (6) implies $\frac{1}{2}q_{k+1}^{-1} > q_k^{-1-\varepsilon}$, so the intervals $(q_k^{-1-\varepsilon}, \frac{1}{2}q_k^{-1})$ cover the interval $(0, 1/2q_1)$. Pick an index m for which $\delta \in (q_m^{-1-\varepsilon}, \frac{1}{2}q_m^{-1})$. For every integer $j > -m/4r$ denote $a_j = q_{m+4rj}$ and set $a_j = 0$ for $j \leq -m/4r$. We have $\delta \in (a_0^{-1-\varepsilon}, \frac{1}{2}a_0^{-1})$, so

$$\delta^{-1/(1+\varepsilon)} < a_0 < \frac{1}{2}\delta^{-1} ,$$

and, by (6),

$$a_{j+1} > a_j^{(1+\varepsilon/2)4r} > a_j^{1+2r\varepsilon} = a_j^{2+2\varepsilon} .$$

It follows that $a_{j+1} \geq 10^{2r} a_j$ for all j , and

$$\begin{aligned} a_1 &> a_0^{2+2\varepsilon} > \delta^{-(2+2\varepsilon)/(1+\varepsilon)} = \delta^{-2} , \\ a_{-1} &< a_0^{1/(2+2\varepsilon)} < a_0^{1/2} < \delta^{-1/2} . \end{aligned}$$

The m th coordinate of the vector $\gamma(y) - \gamma(x)$ can be written as

$$\sum_{j \in \mathbf{Z}} a_j^{-1/2r} E(a_j x) (E(a_j \delta) - 1) .$$

So

$$|\gamma(y) - \gamma(x)| \geq a_0^{-1/2r} |E(a_0 \delta) - 1| - \sum_{j \neq 0} a_j^{-1/2r} |E(a_j \delta) - 1|$$

Since $|E(t) - 1| \geq 4t$ for $0 \leq t \leq 1/2$, we have

$$a_0^{-1/2r} |E(a_0 \delta) - 1| \geq 4\delta a_0^{1-1/2r} > 4\delta^{1-(1-1/2r)/(1+\varepsilon)} = 4\delta^{1/r} .$$

For $j > 0$ we have $a_j > 10^{2r(j-1)} a_1 > 10^{2r(j-1)} \delta^{-2}$, so

$$\begin{aligned} \sum_{j>0} a_j^{-1/2r} |E(a_j \delta) - 1| &\leq \sum_{j>0} 2a_j^{-1/2r} < \sum_{j>0} 2(10^{2r(j-1)} \delta^{-2})^{-1/2r} \\ &= \sum_{j>0} 2 \cdot 10^{-j+1} \delta^{1/r} < 2.5 \delta^{1/r} . \end{aligned}$$

For $-m/4 < j < 0$ we have $10^{4r} < a_j < 10^{j+1}a_{-1} < 10^{j+1}\delta^{-1/2}$, so

$$\begin{aligned} \sum_{j<0} a_j^{-1/2r} |E(a_j\delta) - 1| &\leq \sum_{j<0} 10^{-2} \cdot 2\pi a_j \delta \leq \sum_{j<0} 10^{-1} (10^{j+1}\delta^{-1/2}) \delta \\ &= \sum_{j<0} 10^j \delta^{1/2} < 0.5 \delta^{1/2} \leq 0.5 \delta^{1/r}. \end{aligned}$$

It follows that $|\gamma(x) - \gamma(y)| \leq (4 - 2.5 - 0.5)\delta^{1/r} = \delta^{1/r}$. \square

4.3. Proposition. *For every integer $n \geq 7$ and $v \in \mathbf{R}^n$ there is a three-dimensional smooth submanifold $M \subset T^n$ and a smooth flow on M , one of whose trajectories has the rotation vector v (as a curve in T^n).*

Proof. Let M_0 be a two-dimensional orientable surface of genus n equipped with a negatively curved Riemannian metric, and let UTM_0 be its unit tangent bundle. Every map $UTM_0 \rightarrow T^n$ is homotopic to a self-transversal smooth map which has to be an embedding since $n \geq 7$. Choose such an embedding $UTM_0 \hookrightarrow T^n$ which induces an epimorphism of fundamental groups; let M be the image of UTM_0 in T^n .

We consider flows on $M \simeq UTM_0$ preserving the Liouville measure. Each flow is generated by a divergence-free vector field on M . The *rotation vector* $\mathcal{R}(X) \in \mathbf{R}^n \cong H_1(T^n; \mathbf{R})$ of such a vector field X is the average value of rotation vectors of its trajectories as curves in T^n . The map $X \rightarrow \mathcal{R}(X)$ is linear and the set of its values is the entire \mathbf{R}^n . (To prove this, observe that a homology class of any closed geodesic in UTM_0 can be represented as a rotation vector of some flow concentrated in a small neighborhood of the geodesic.) Let Y be a divergence-free vector field with $\mathcal{R}(Y) = v$.

Let X_0 be a vector field which generates the geodesic flow of our metric. It is skew-symmetric with respect to the reflection $w \mapsto -w$ of UTM_0 , so $\mathcal{R}(X_0) = 0$. Since the metric has negative curvature, its geodesic flow is an Anosov's flow. Hence the vector field $X_\varepsilon := X_0 + \varepsilon Y$ still generates ergodic flow for ε small enough. In particular, almost every trajectory generated by X_ε has the rotation vector equal to $\mathcal{R}(X_\varepsilon) = \varepsilon v$. To complete the proof, consider the flow generated by $X_\varepsilon/\varepsilon$. \square

4.4. Let $n = 8r + 6$, $v \in \mathbf{R}^n$ be a completely irrational vector. Fix $M \subset T^n$ and a trajectory $\gamma: \mathbf{R} \rightarrow M$ of a smooth flow on M constructed by Proposition 4.3. For each $x \in T^n$, identify the tangent space $T_x T^n$ with \mathbf{R}^n in the standard way, and then consider the family $\{L_x = T_x M : x \in M\}$ of 3-dimensional linear subspaces of \mathbf{R}^n . Pick a $(8r)$ -dimensional linear subspace $L \subset \mathbf{R}^n$ which is transversal to this family, i. e. $L \cap L_x = \{0\}$ for all $x \in M$. Transversality implies that there exist $\varepsilon > 0$, a tubular neighborhood $U \subset T^n$ of M , and a smooth retraction $p: U \rightarrow M$ such that $p(x+y) = x$ whenever $x \in M$, $y \in L$ and $|y| < \varepsilon$.

Let $\alpha \in \mathbf{R}$ be an approximable number. Using Proposition 4.2, construct an r -stretching 1-periodic continuous curve $\gamma_1: \mathbf{R} \rightarrow L$ such that the map $f: \mathbf{R} \rightarrow L$ given by

$$f(s) = \gamma_1(s + \alpha) - \gamma_1(s)$$

is C^1 -smooth. After a suitable homothety we may assume that $|\gamma_1(s)| < \varepsilon/8$, and hence $|f(s)| < \varepsilon/4$, for all $s \in \mathbf{R}$.

Denote $\sigma = 1/3$. Fix a smooth function $\varphi: [0, 1 + \sigma] \rightarrow [0, 1]$ such that $\varphi(t) = 0$ for $t \in [0, \sigma]$ and $\varphi(t) = 1$ for $t \in [1, 1 + \sigma]$. Define $\Gamma: \mathbf{R} \times [0, 1 + \sigma] \rightarrow T^{n+1} \times \mathbf{R}$ by

$$\Gamma(s, t) = (\gamma(s + \alpha t) + \gamma_1(s) + f(s)\varphi(t), (s + \alpha t) \bmod 1, t) \in T^n \times S^1 \times \mathbf{R}.$$

4.5. Lemma. *There exists $c > 0$ such that*

$$(7) \quad |\Gamma(s, t) - \Gamma(s', t')| \geq c |(s - s') \bmod 1|^{1/r}$$

for all $s, s' \in \mathbf{R}$, $t, t' \in [0, 1 + \sigma]$.

Proof. For values of $|(s - s') \bmod 1|$ bounded away from zero, we may find a constant c satisfying (7) since

$$|\Gamma(s, t) - \gamma_2(s', t')| \geq \min\{1, 1/|\alpha|\} \cdot |(s - s') \bmod 1|$$

and the function $x \mapsto x^{1/r}/x$ is bounded outside any neighborhood of zero. So it suffices to prove the statement of lemma when $|(s - s') \bmod 1|$ is sufficiently small.

First let $t' = t = 0$. We have

$$\begin{aligned} |\Gamma(s, 0) - \Gamma(s', 0)| &\geq |\gamma(s) + \gamma_1(s) - \gamma_1(s') - \gamma(s' + \alpha t)| \\ &\geq \text{dist}(M + \gamma_1(s) - \gamma_1(s'), M) \geq c_1 |\gamma_1(s) - \gamma_1(s')| \end{aligned}$$

since the vector $\gamma_1(s) - \gamma_1(s')$ lies in L and its length is less than ε . Then (7) for $t' = t = 0$ follows from the fact that γ_1 is r -stretching and 1-periodic.

Since $\Gamma(s, t) - \Gamma(s, 0)$ is a C^1 (and hence Lipschitz) function of (s, t) , the estimate (7) for $t' = t = 0$ implies the same for $t' = t \neq 0$. Then the complete statement follows since $|t - t'| \leq |\Gamma(s, t) - \Gamma(s, t')| \leq C \cdot |t - t'|$ for some constant C . \square

Define $\Gamma_0: \mathbf{R} \times [0, 1 + \sigma] \rightarrow T^{n+2}$ to be $\pi \circ \Gamma$ where $\pi: T^{n+1} \times \mathbf{R} \rightarrow T^{n+1} \times S^1 = T^{n+1}$ is the standard factorization. Note that for $t \in [0, \sigma]$,

$$\begin{aligned} \Gamma(s, t + 1) &= (\gamma(s + \alpha + \alpha t) + \gamma_1(s) + f(s), (s + \alpha + \alpha t) \bmod 1, t + 1) \\ &= \Gamma(s + \alpha, t) + (0, 0, 1) \in T^n \times S^1 \times \mathbf{R}, \end{aligned}$$

so

$$(8) \quad \Gamma_0(s, t + 1) = \Gamma_0(s + \alpha, t) \quad \text{for all } s \in \mathbf{R}, t \in [0, \sigma].$$

In other words, a piece of the surface $\Gamma_0(s, t)$ where $t \in [1, 1 + \sigma]$ matches the one where $t \in [0, \sigma]$, up to the parameter shift $s \mapsto s + \alpha$. Γ has no other self-intersections since the last coordinate of $\Gamma_0(s, t)$ is $t \bmod 1$.

Let w_0 be the vector field along Γ_0 given by $w_0(s, t) = \frac{d}{dt} \Gamma_0(s, t)$. We are going to consider $w_0(s, t)$ as a function of point $\Gamma_0(s, t) \in T^{n+2}$. Though $w_0(s, t)$ is only C^1 as a function of (s, t) , it turns out to be smoother when viewed as a function on a subset of T^{n+2} .

4.6. Lemma. *There exists a C^{r-1} -smooth vector field W_0 on T^{n+2} such that $w_0 = W_0 \circ \Gamma_0$.*

Proof. In view of (8) it suffices to prove the same statement for a similar vector field along Γ , $w(s, t) = \frac{d}{dt} \Gamma(s, t)$. By Whitney's extension theorem [?] it suffices to construct for each $(s, t) \in \mathbf{R} \times [0, 1 + \sigma]$ a smooth vector field $\varphi_{s,t}$ in a neighborhood of $\Gamma(s, t) \in T^{n+1} \times \mathbf{R}$ so that

$$|w(\Gamma(s', t')) - \varphi_{s,t}(s', t')| \leq O(|\Gamma(s', t') - \Gamma(s, t)|^r), \quad (s', t') \rightarrow (s, t).$$

Recall that γ is a trajectory of a smooth flow on $M \subset T^n$, so there is a smooth vector field V on M such that $\gamma'(t) = V(\gamma(t))$ for all $t \in \mathbf{R}$. So

$$w(s, t) = (\alpha V(\gamma(s + \alpha t)) + f(s)g'(t), \alpha, 1)$$

is a C^1 -smooth function of (s, t) . In particular,

$$(9) \quad |w(t', \tau') - w(t, \tau')| \leq C|t' - t|$$

for some $C > 0$. Let U be a neighborhood of $M \subset T^n$ and $p: U \rightarrow M$ be a retraction as described in 4.4. Define $\varphi_{s,t}$ by

$$\begin{aligned} \varphi_{s,t}(x, y, z) &= (\alpha V(p(x - f(s)g(z))) + f(s)g'(z), \alpha, 1), \\ x &\in U + f(s)g(z) \subset T^n, \quad y \in S^1, \quad z \in [0, 1 + \sigma]. \end{aligned}$$

Then $\varphi_{s,t}$ is a smooth function and $\varphi_{s,t}(\Gamma(s, t)) = w(s, t)$ for all $s \in \mathbf{R}$, $t \in [0, 1 + \sigma]$. Since f is 1-periodic, $\varphi_{s+m,t} = \varphi_{s,t}$ for any $m \in \mathbf{Z}$. Also, $\varphi_{s,t}$ does not depend on t and $\varphi_{s,t}(x, y, z)$ is a C^1 -smooth function of (s, x, y, z) . Then for any $m \in \mathbf{Z}$ we have

$$|w(s', t') - \varphi_{s,t}(\Gamma(s', t'))| = |\varphi_{s',t'}(\Gamma(s', t')) - \varphi_{s+m,t'}(\Gamma(s', t'))| \leq C|s' - s - m|$$

for some constant C . Since m is arbitrary,

$$|w(s', t') - \varphi_{s,t}(\Gamma(s', t'))| \leq C|(s' - s) \bmod 1|$$

Using 4.5 we conclude that

$$|w(s', t') - \varphi_{s,t}(\Gamma(s', t'))| \leq C|\Gamma(s', t') - \Gamma(s, t)|^r,$$

and the lemma follows. \square

Thus we have a flow on T^{n+2} (generated by a C^{r-1} -smooth vector field W_0) such that all the curves $\Gamma_0^s := \Gamma_0(s, \cdot): [0, 1] \rightarrow T^{n+2}$ are intervals of its trajectories. Every curve Γ_0^s joins the points $(\gamma(s), s, 0)$ and $(\gamma(s + \alpha), s + \alpha, 0)$ in T^{n+2} , and then the similar curve $\Gamma_0^{s+\alpha}$ forms the next piece of a trajectory. Clearly Γ_0^s is homotopic to a curve $t \mapsto (\gamma(s + \alpha t), s + \alpha t, t) \in T^{n+2}$, so the rotation vector of the entire trajectory is equal to $(\alpha v, \alpha, 1)$, where $v \in \mathbf{R}^n$ is the rotation vector of γ (see 4.4).

Since the last two coordinates of $w_0(\cdot)$ are the constants α and 1, we may assume the same for $W_0(\cdot)$, i.e., $W_0(\cdot) = (W_1(\cdot), \alpha, 1)$ for some smooth $W_1: T^{n+2} \rightarrow \mathbf{R}^n$.

4.7. Let $\mathcal{M} \subset T^{n+2}$ be the image of Γ_0 . Let \tilde{W}_0 and $\tilde{\mathcal{M}}$ be the lifts of W_0 and \mathcal{M} from T^{n+2} to \mathbf{R}^{n+2} . Both \tilde{W}_0 and $\tilde{\mathcal{M}}$ are \mathbf{Z}^{n+2} -invariant. Define two linear functions L and H on \mathbf{R}^{n+2} by $L(x, y, z) = y - \alpha z$ and $H(x, y, z) = z$ for $x \in \mathbf{R}^n$, $y, z \in \mathbf{R}$. Here are the properties of \tilde{W}_0 and $\tilde{\mathcal{M}}$ that we will need:

- (1) \tilde{W}_0 is a C^{r-1} -smooth vector field whose last two coordinates are α and 1. In particular, L is constant and H increases at the constant rate 1 under the flow generated by \tilde{W}_0 .
- (2) $\tilde{\mathcal{M}} \subset \mathbf{R}^{n+2}$ is invariant under the flow generated by \tilde{W}_0 . At least one trajectory of \tilde{W}_0 lies in $\tilde{\mathcal{M}}$ and has rotation vector $(\alpha v, \alpha, 1)$.
- (3) There exists $c > 0$ such that

$$|p - q| \geq c|L(p) - L(q)|^{1/r} \quad \text{for all } x, y \in \tilde{\mathcal{M}}.$$

These properties follow immediately from Lemma 4.5.

Let g_0 be a Riemannian metric on \mathbf{R}^{n+2} which induces the standard flat metric on every hyperplane $\mathbf{R}^{n+1} \times \{\text{const}\}$ and such that \tilde{W}_0 is a unit-length vector field orthogonal to these hyperplanes with respect to g_0 . Clearly g_0 is uniquely determined, C^{r-1} -smooth, and \mathbf{Z}^{n+2} -periodic. Let ρ_0 denote the distance associated with g_0 . Note that L and H are 1-Lipschitz functions with respect to ρ_0 .

It is easy to construct a C^{r-2} -smooth \mathbf{Z}^n -periodic function $\varphi: \mathbf{R}^{n+2} \rightarrow \mathbf{R}$ such that $\varphi|_{\tilde{\mathcal{M}}} \equiv 0$ and $\varphi(x) \geq \rho_0(x, \tilde{\mathcal{M}})^{r-1}$ for all $x \in \mathbf{R}^{n+2}$. Define a Riemannian structure g on \mathbf{R}^{n+2} by $g = g_0/(1 + \varphi)^2$. g_0 is periodic and C^{r-2} -smooth.

Let ρ be the metric determined by g and let $\|\cdot\|$ be the stable norm of ρ . We will prove that the unit sphere of $\|\cdot\|$ is nonsmooth at the point $(\alpha v, \alpha, 1) \in \mathbf{R}^{n+2}$. First note that $\|(\alpha v, \alpha, 1)\| \leq 1$ by (2) above.

4.8. Lemma. *There exists an $\varepsilon > 0$ such that*

$$\rho(p, q) \geq |H(p) - H(q)| + \varepsilon|L(p) - L(q)|$$

whenever $H(p) - H(q)$ is integer, $p, q \in \mathbf{R}^{n+2}$.

Proof. Without loss of generality we may assume $0 \leq H(p) < 1$ and $H(q) \geq H(p)$. Since ρ is a length metric, there is a sequence of points $p_0 = p, p_1, \dots, p_N = q$ such that $\rho(p, q) = \sum \rho(p_i, p_{i+1})$ and for each $i = 0, \dots, N-1$ either $H(p_{i+1}) = H(p_i)$, or $H(p_{i+1}) = H(p_i) + 1$ and a shortest curve joining p_i and p_{i+1} lies between the hyperplanes $H^{-1}(H(p_i))$ and $H^{-1}(H(p_{i+1}))$. It suffices to prove the desired estimate for each pair of points (p_i, p_{i+1}) instead of (p, q) . In the case $H(p_i) = H(p_{i+1})$ it holds with $\varepsilon = 1$ because L is 1-Lipschitz with respect to ρ .

Let $H(q) = H(p) + 1$ and let γ be a C^1 -smooth curve joining p and q and which lies between the hyperplanes $\{H = H(p)\}$ and $\{H = H(q)\}$. We assume that $\gamma: [0, 1] \rightarrow (\mathbf{R}^{n+2}, \rho)$ is a constant speed parametrization. Let $\ell(\gamma)$ denote the length of γ in (\mathbf{R}^{n+2}, ρ) . We have to check that $\ell(\gamma) \geq 1 + \varepsilon|L(p) - L(q)|$ for some $\varepsilon > 0$ independent of γ, p and q .

Consider the map $f: \mathbf{R}^{n+1} \times [0, 2] \rightarrow \mathbf{R}^{n+2}$ defined as follows: for $x \in \mathbf{R}^{n+1}$, $y \in [0, 2]$, $f(x, y)$ is the y -shift of the point $(x, 0) \in \mathbf{R}^{n+2}$ along the vector field \tilde{W}_0 . Clearly f preserves hyperplanes of the form $\{H = \text{const}\}$, and f^{-1} is a change of variables in $\mathbf{R}^{n+1} \times [0, 2]$ which transforms \tilde{W}_0 into the $(n+2)^{\text{nd}}$ coordinate vector field. Since f is a \mathbf{Z}^{n+1} -periodic C^1 -smooth diffeomorphism, there are constants $c_1, c_2 > 0$ such that

$$|x - y|/c_1 \geq \rho_0(f(x), f(y)) \geq c_1|x - y| \quad \text{for all } x, y \in \mathbf{R}^{n+1} \times [0, 2]$$

and

$$|x - y| \geq c_2|L \circ f(x) - L \circ f(y)|^{1/r} \quad \text{whenever } x, y \in f^{-1}(\tilde{\mathcal{M}}).$$

Note that $L \circ f$ is the $(n+1)^{\text{st}}$ coordinate function on $\mathbf{R}^{n+1} \times [0, 2]$.

We may write $\gamma(t) = f(\gamma_1(t), \gamma_2(t))$ for some $\gamma_1: [0, 1] \rightarrow \mathbf{R}^{n+1}$ and $\gamma_2: [0, 1] \rightarrow [0, 2]$. Clearly $g_2(t) = H \circ \gamma(t)$ for all t . Since the vector field \tilde{W}_0 is orthogonal to the hyperplanes $\{H = \text{const}\}$ with respect to \tilde{g} , we have

$$\|\gamma'(t)\|_g = (1 + \varphi(\gamma(t))) \cdot \|g'(t)\|_{g_0} \geq (1 + \varphi(\gamma(t))) \sqrt{c_1 \gamma_1'^2 + \gamma_2'^2}.$$

Let $\delta = |L(p) - L(q)|$. Consider the following cases.

Case 1: $\delta \geq 2$. Then $\ell(\gamma) \geq \rho(p, q) \geq |L(p) - L(q)| = \delta \geq 1 + \delta/2$. In the remaining cases we suppose $\delta < 2$.

Case 2: $\rho_0(p, \tilde{\mathcal{M}})$ and $\rho_0(q, \tilde{\mathcal{M}})$ are less than $c_3\delta^{1/r}$ where $c_3 = c_1c_2/3$. It follows that $\gamma_1(0)$ and $\gamma_1(1)$ lie within $(\frac{1}{3}c_2\delta^{1/r})$ -neighborhood of $f^{-1}(\tilde{\mathcal{M}})$ in $\mathbf{R}^{n+1} \times [0, 2]$, so $|\gamma_1(0) - \gamma_1(1)| \geq \frac{1}{3}c_2\delta^{1/r}$. Then

$$\begin{aligned} \ell(\gamma)^2 &= \int_0^1 \|\gamma'(t)\|_g^2 \geq \int_0^1 c_1\gamma_1'^2 + \gamma_2'^2 \\ &\geq c_1|\gamma_1(1) - \gamma_1(0)|^2 + |\gamma_2(1) - \gamma_2(0)|^2 \geq 1 + \frac{c_1c_2^2}{9}\delta^{2/r}. \end{aligned}$$

Since $r \geq 2$, it follows that $\ell(\gamma) \geq 1 + c_4\delta$ where c_4 can be easily expressed in terms of c_1, c_2 and the upper bound for δ (that is, 2).

Case 3: $\rho_0(p, \tilde{\mathcal{M}}) \geq c_3\delta^{1/r}$ (or similar case with q). Let ℓ_0 denote length with respect to ρ_0 and let γ_0 be the starting interval of γ such that $\ell_0(\gamma_0) = \frac{1}{2}c_3\delta^{1/r}$. Then γ_0 lies entirely outside the $(\frac{1}{2}c_3\delta^{1/r})$ -neighborhood of \mathcal{M} in $(\mathbf{R}^{n+2}, \rho_0)$. In this region, we have $\varphi \geq (\frac{1}{2}c_3\delta^{1/r})^{r-1} = c_5\delta^{1-1/r}$. Therefore

$$\ell(\gamma) \geq \ell_0(\gamma) + c_5\delta^{1-1/r}\ell_0(\gamma_0) \geq 1 + c_6\delta^{1/r}\delta^{1-1/r} = 1 + c_6\delta$$

where $c_6 = c_3c_5/2$.

In all three cases, $\ell(\gamma) \geq 1 + \varepsilon\delta$ for $\varepsilon = \min\{1/2, c_4, c_6\}$. \square

4.9. Corollary. $\|\cdot\| \geq H + \varepsilon|L|$. \square

Since $\|(\alpha v, \alpha, 1)\| = 1$ and $L(\alpha v, \alpha, 1) = 0$, both of the linear functions $H - \varepsilon L$ and $H + \varepsilon L$ support the unit ball of $\|\cdot\|$ at the point $(\alpha v, \alpha, 1)$. It follows that the unit sphere of $\|\cdot\|$ is nonsmooth at $(\alpha v, \alpha, 1)$.

REFERENCES

- [Ba1] V. Bangert, *Minimal Geodesics*, Erg. Theory and Dyn. Syst. **10**, 263–286.
- [Ba2] V. Bangert, *Geodesic rays, Busemann functions and monotone twist maps*, Calc. Var. **2**, 49–63.
- [Ba3] V. Bangert, *Mather sets for twist maps and geodesics on tori*, In: Dynamics reported **1**, 1–56.
- [Ba4] V. Bangert, *Question*, In: Address to Zurich IMC.
- [Bu1] D. Burago, *Periodic metrics*, Advances in Soviet Math. **9** (1992), New York, 205–210.
- [Bu2] ———, *Periodic metrics*, in “Seminar on Dynamical Systems”, Progress in Nonlinear Differential Equations (H. Brezis, ed.), vol. 12, Birkhauser, 1994, pp. 90–96.
- [BuI] D. Burago and S. Ivanov, *Riemannian tori without conjugate points are flat*, GAFA **4:3** (1994), 259–269.
- [BuI2] D. Burago and S. Ivanov, *On asymptotic volume of tori*, GAFA **5:5** (1995), 800–808.
- [BuI3] D. Burago and S. Ivanov, *Isometric embeddings of Finsler manifolds*, St. Petersburg Math. Journal **5:1** (1994), 159–169.
- [CK] C. Croke, B. Kleiner, *On tori without conjugate points*, Invent. Math. **120** (1995), 241–257.
- [G] M. Gromov, *Asymptotic Invariants of Infinite Groups*, Geometric Group Theory vol 2, London Math. Society Lecture Notes 182 (1993).
- [GLP] M. Gromov, J. Lafontaine, P. Pansu, *Structure metriques pour les varietés Riemanniennes*, CEDIC\Fernand Math., Paris.
- [H] Jens Heber, *On the geodesic flow of tori without conjugate points*, Math. Z. **216** (1994), 209–216.
- [Ka] M. Kac, *Statistical independance in probability*, Analysis and Number Theory, Carus Math. Monograph 12, Math. Assoc. Amer (1959).

- [M1] John Mather, *Variational construction of connecting orbits*, Ann. Inst. Fourier (Grenoble) **43** (1993), no. 5, 1349-1386.
- [M2] ———, *Action minimizing invariant measures for positive definite Lagrangian systems*, Math. Z. **207** (1991), no. 2, 169-207.
- [M3] ———, *Minimal measures*, Comm. Math. Helv. **64**, 375-394.
- [M4] ———, *Differentiability of the minimal average action as a function of the rotation number*, Bull. Soc. Bras. Mat. **21**, 59-70.
- [Na] Fedor Nazarov, Lecture given at Steklov institute, St. Petersburg.

DMITRI BURAGO, DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16801

E-mail address: burago@math.psu.edu

SERGEI IVANOV, STEKLOV INSTITUTE OF MATHEMATICS, 191011, FONTANKA 27, ST. PETERSBURG, RUSSIA

E-mail address: svivanov@pdmi.ras.ru

BRUCE KLEINER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA 19104-6395

E-mail address: bkleiner@math.upenn.edu